



INTERNATIONAL APPLICATION PUBLISHED UNDER THE PATENT COOPERATION TREATY (PCT)

<p>(51) International Patent Classification ⁶ : C07D 253/075, 401/12, 403/12, 413/10, 417/10, 495/04, 409/10, 413/12, 417/14, A61K 31/53, C07D 401/14, 413/14 // (C07D 495/04, 213:00, 333:00)</p>	<p>A1</p>	<p>(11) International Publication Number: WO 99/02505 (43) International Publication Date: 21 January 1999 (21.01.99)</p>
<p>(21) International Application Number: PCT/EP98/04191 (22) International Filing Date: 7 July 1998 (07.07.98) (30) Priority Data: 97202118.2 10 July 1997 (10.07.97) EP (71) Applicant (for all designated States except US): JANSSEN PHARMACEUTICA N.V. [BE/BE]; Turnhoutseweg 30, B-2340 Beerse (BE). (72) Inventors; and (75) Inventors/Applicants (for US only): LACRAMPE, Jean, Fernand, Armand [FR/FR]; Janssen-Cilag S.A., 1, rue Camille Desmoulins, TSA 91003, F-92787 Issy-les-Moulineaux Cedex 9 (FR). FREYNE, Eddy, Jean, Edgard [BE/BE]; Janssen Pharmaceutica N.V., Turnhout- seweg 30, B-2340 Beerse (BE). VENET, Marc, Gaston [FR/FR]; Janssen-Cilag S.A., 1, rue Camille Desmoulins, TSA 91003, F-92787 Issy-les-Moulineaux Cedex 9 (FR). BOECKX, Gustaaf, Maria [BE/BE]; Janssen Pharmaceutica N.V., Turnhoutseweg 30, B-2340 Beerse (BE).</p>		<p>(81) Designated States: AL, AM, AT, AU, AZ, BA, BB, BG, BR, BY, CA, CH, CN, CU, CZ, DE, DK, EE, ES, FI, GB, GE, GH, GM, GW, HR, HU, ID, IL, IS, JP, KE, KG, KP, KR, KZ, LC, LK, LR, LS, LT, LU, LV, MD, MG, MK, MN, MW, MX, NO, NZ, PL, PT, RO, RU, SD, SE, SG, SI, SK, SL, TJ, TM, TR, TT, UA, UG, US, UZ, VN, YU, ZW, ARIPO patent (GH, GM, KE, LS, MW, SD, SZ, UG, ZW), Eurasian patent (AM, AZ, BY, KG, KZ, MD, RU, TJ, TM), European patent (AT, BE, CH, CY, DE, DK, ES, FI, FR, GB, GR, IE, IT, LU, MC, NL, PT, SE), OAPI patent (BF, BJ, CF, CG, CI, CM, GA, GN, ML, MR, NE, SN, TD, TG).</p> <p>Published With international search report. Before the expiration of the time limit for amending the claims and to be republished in the event of the receipt of amendments.</p>
<p>(54) Title: IL-5 INHIBITING 6-AZAUACIL DERIVATIVES</p> <div data-bbox="631 1123 1031 1291" data-label="Chemical-Block"> </div> <p>(57) Abstract</p> <p>The present invention is concerned with the compounds of formula (I) the <i>N</i>-oxides, the pharmaceutically acceptable addition salts and the stereochemically isomeric forms thereof, p and q are 0, 1, 2, 3 or 4 and q is also 5; X is O, S, NR³ or a direct bond; R¹ is hydrogen, hydroxy, halo, optionally substituted amino, optionally substituted C₁₋₆alkyl, C₁₋₆alkyloxy, C₃₋₇cycloalkyl or aryl; R² is aryl, Het¹, C₃₋₇cycloalkyl, optionally substituted C₁₋₆alkyl; and if X is O, S or NR³, then R² may also be a carbonyl or thiocarbonyl linked substituent; R³ is hydrogen or C₁₋₄alkyl; R⁴ and R⁵ independently are optionally substituted C₁₋₆alkyl, halo, hydroxy, mercapto, C₁₋₆alkyloxy, C₁₋₆alkylthio, C₁₋₆alkylcarbonyloxy, aryl, cyano, nitro, Het³, R⁶ or NR⁷R⁸; R⁶ is substituted sulfonyl or sulfinyl; R⁷ and R⁸ are hydrogen, optionally substituted C₁₋₄alkyl, aryl, a carbonyl or thiocarbonyl linked substituent, C₃₋₇cycloalkyl, Het³ and R⁶; R⁹ and R¹⁰ are each independently selected from hydrogen, optionally substituted C₁₋₄alkyl, phenyl, a carbonyl or thiocarbonyl linked substituent, C₃₋₇cycloalkyl, Het³ and R⁶; R¹¹ is hydroxy, mercapto, cyano, nitro, halo, trihalomethyl, C₁₋₄alkyloxy, carboxyl, C₁₋₄alkyloxycarbonyl, trihaloC₁₋₄alkylsulfonyloxy, R⁶, NR⁷R⁸, C(=O)NR⁷R⁸, aryl, aryloxy, arylcarbonyl, C₃₋₇cycloalkyl, C₃₋₇cycloalkyloxy, phthalimide-2-yl, Het³ and C(=O)Het³; R¹² and R¹³ are each independently selected from hydrogen, optionally substituted C₁₋₄alkyl, phenyl, a carbonyl or thiocarbonyl linked substituent, C₃₋₇cycloalkyl and R⁶; aryl is optionally substituted phenyl; Het¹, Het² and Het³ are optionally substituted heterocycles; to processes for their preparation and compositions comprising them. It further relates to their use as a medicine.</p>		

FOR THE PURPOSES OF INFORMATION ONLY

Codes used to identify States party to the PCT on the front pages of pamphlets publishing international applications under the PCT.

AL	Albania	ES	Spain	LS	Lesotho	SI	Slovenia
AM	Armenia	FI	Finland	LT	Lithuania	SK	Slovakia
AT	Austria	FR	France	LU	Luxembourg	SN	Senegal
AU	Australia	GA	Gabon	LV	Latvia	SZ	Swaziland
AZ	Azerbaijan	GB	United Kingdom	MC	Monaco	TD	Chad
BA	Bosnia and Herzegovina	GE	Georgia	MD	Republic of Moldova	TG	Togo
BB	Barbados	GH	Ghana	MG	Madagascar	TJ	Tajikistan
BE	Belgium	GN	Guinea	MK	The former Yugoslav Republic of Macedonia	TM	Turkmenistan
BF	Burkina Faso	GR	Greece	ML	Mali	TR	Turkey
BG	Bulgaria	HU	Hungary	MN	Mongolia	TT	Trinidad and Tobago
BJ	Benin	IE	Ireland	MR	Mauritania	UA	Ukraine
BR	Brazil	IL	Israel	MW	Malawi	UG	Uganda
BY	Belarus	IS	Iceland	MX	Mexico	US	United States of America
CA	Canada	IT	Italy	NE	Niger	UZ	Uzbekistan
CF	Central African Republic	JP	Japan	NL	Netherlands	VN	Viet Nam
CG	Congo	KE	Kenya	NO	Norway	YU	Yugoslavia
CH	Switzerland	KG	Kyrgyzstan	NZ	New Zealand	ZW	Zimbabwe
CI	Côte d'Ivoire	KP	Democratic People's Republic of Korea	PL	Poland		
CM	Cameroon	KR	Republic of Korea	PT	Portugal		
CN	China	KZ	Kazakhstan	RO	Romania		
CU	Cuba	LC	Saint Lucia	RU	Russian Federation		
CZ	Czech Republic	LI	Liechtenstein	SD	Sudan		
DE	Germany	LK	Sri Lanka	SE	Sweden		
DK	Denmark	LR	Liberia	SG	Singapore		
EE	Estonia						

IL-5 inhibiting 6-Azaauracil derivatives


The present invention concerns novel IL-5 inhibiting 6-azauracil derivatives useful for
 5 treating eosinophil-dependent inflammatory diseases; to processes for their preparation
 and compositions comprising them. It further relates to their use as a medicine.

Eosinophil influx, leading to subsequent tissue damage, is an important pathogenic event
 in bronchial asthma and allergic diseases. The cytokine interleukin-5 (IL-5), produced
 10 mainly by T lymphocytes as a glycoprotein, induces the differentiation of eosinophils in
 bone marrow and, primes eosinophils for activation in peripheral blood and sustains their
 survival in tissues. As such, IL-5 plays a critical role in the process of eosinophilic
 inflammation. Hence, the possibility that inhibitors of IL-5 production would reduce the
 production, activation and/or survival of eosinophils provides a therapeutic approach to
 15 the treatment of bronchial asthma and allergic diseases such as, atopic dermatitis,
 allergic rhinitis, allergic conjunctivitis, and also other eosinophil-dependent inflammatory
 diseases.

Steroids, which strongly inhibit IL-5 production *in vitro*, have long been used as the
 20 only drugs with remarkable efficacy for bronchial asthma and atopic dermatitis, but they
 cause various serious adverse reactions such as diabetes, hypertension and cataracts.
 Therefore, it would be desirable to find non-steroidal compounds having the ability to
 inhibit IL-5 production in human T-cells and which have little or no adverse reactions.

US 4,631,278 discloses α -aryl-4-(4,5-dihydro-3,5-dioxo-1,2,4-triazin-2(3H)-
 25 yl)benzeneacetonitriles and US 4,767,760 discloses 2-(substituted phenyl)-1,2,4-
 triazine-3,5(2H,4H)-diones, all having anti-protozoal activity, in particular, anti-
 coccidial activity. EP 831,088 discloses 1,2,4-triazine-3,5-diones as anticoccidial agents.
 Unexpectedly, the 6-azauracil derivatives of the present invention prove to be potent
 30 inhibitors of the production of IL-5.

The present invention is concerned with the compounds of formula



the *N*-oxides, the pharmaceutically acceptable addition salts and the stereochemically

isomeric forms thereof, wherein :

p represents an integer being 0, 1, 2, 3 or 4;

q represents an integer being 0, 1, 2, 3, 4 or 5;

X represents O, S, NR³ or a direct bond;

5 R¹ represents hydrogen, hydroxy, halo, amino, mono- or di(C₁₋₄alkyl)amino, C₁₋₆alkyl, C₁₋₆alkyloxy, C₃₋₇cycloalkyl, aryl, arylC₁₋₆alkyl, aminoC₁₋₄alkyl, mono- or di(C₁₋₄alkyl)aminoC₁₋₄alkyl or mono- or di(C₁₋₄alkyl)aminoC₁₋₄alkylamino;

10 R² represents aryl, Het¹, C₃₋₇cycloalkyl, C₁₋₆alkyl or C₁₋₆alkyl substituted with one or two substituents selected from hydroxy, cyano, amino, mono- or di(C₁₋₄alkyl)amino, C₁₋₆alkyloxy, C₁₋₆alkylsulfonyloxy, C₁₋₆alkyloxycarbonyl, C₃₋₇cycloalkyl, aryl, aryloxy, arylthio, Het¹, Het¹oxy and Het¹thio; and if X is O, S or NR³, then R² may also represent aminocarbonyl, aminothiocarbonyl, C₁₋₄alkylcarbonyl, C₁₋₄alkylthiocarbonyl, arylcarbonyl or arylthiocarbonyl;

R³ represents hydrogen or C₁₋₄alkyl;

15 each R⁴ independently represents C₁₋₆alkyl, halo, polyhaloC₁₋₆alkyl, hydroxy, mercapto, C₁₋₆alkyloxy, C₁₋₆alkylthio, C₁₋₆alkylcarbonyloxy, aryl, cyano, nitro, Het³, R⁶, NR⁷R⁸ or C₁₋₄alkyl substituted with Het³, R⁶ or NR⁷R⁸;

each R⁵ independently represents C₁₋₆alkyl, halo, polyhaloC₁₋₆alkyl, hydroxy, mercapto, C₁₋₆alkyloxy, C₁₋₆alkylthio, C₁₋₆alkylcarbonyloxy, aryl, cyano, nitro, Het³, R⁶, NR⁷R⁸ or C₁₋₄alkyl substituted with Het³, R⁶ or NR⁷R⁸;

20 each R⁶ independently represents C₁₋₆alkylsulfonyl, aminosulfonyl, mono- or di-(C₁₋₄alkyl)aminosulfonyl, mono- or di(benzyl)aminosulfonyl, polyhaloC₁₋₆alkylsulfonyl, C₁₋₆alkylsulfinyl, phenylC₁₋₄alkylsulfonyl, piperazinylsulfonyl, aminopiperidinylsulfonyl, piperidinylaminosulfonyl, N-C₁₋₄alkyl-N-piperidinylaminosulfonyl;

25 each R⁷ and each R⁸ are independently selected from hydrogen, C₁₋₄alkyl, hydroxy-C₁₋₄alkyl, dihydroxyC₁₋₄alkyl, aryl, arylC₁₋₄alkyl, C₁₋₄alkyloxyC₁₋₄alkyl, C₁₋₄alkylcarbonyl, arylcarbonyl, C₁₋₄alkylcarbonyloxyC₁₋₄alkylcarbonyl, hydroxyC₁₋₄alkylcarbonyl, C₁₋₄alkyloxycarbonylcarbonyl, mono- or di(C₁₋₄alkyl)aminoC₁₋₄alkyl, arylaminocarbonyl, arylaminothiocarbonyl, Het³aminocarbonyl, Het³aminothiocarbonyl, C₃₋₇cycloalkyl, pyridinylC₁₋₄alkyl, Het³ and R⁶;

30 R⁹ and R¹⁰ are each independently selected from hydrogen, C₁₋₄alkyl, hydroxyC₁₋₄alkyl, dihydroxyC₁₋₄alkyl, phenyl, phenylC₁₋₄alkyl, C₁₋₄alkyloxyC₁₋₄alkyl, C₁₋₄alkylcarbonyl, phenylcarbonyl, C₁₋₄alkylcarbonyloxyC₁₋₄alkylcarbonyl, hydroxyC₁₋₄alkylcarbonyl, C₁₋₄alkyloxycarbonylcarbonyl, mono- or di(C₁₋₄alkyl)aminoC₁₋₄alkyl, 35 phenylaminocarbonyl, phenylaminothiocarbonyl, Het³aminocarbonyl, Het³aminothiocarbonyl, C₃₋₇cycloalkyl, pyridinylC₁₋₄alkyl, Het³ and R⁶;

each R¹¹ independently being selected from hydroxy, mercapto, cyano, nitro, halo,

trihalomethyl, C₁₋₄alkyloxy, carboxyl, C₁₋₄alkyloxycarbonyl,

trihaloC₁₋₄alkylsulfonyloxy, R⁶, NR⁷R⁸, C(=O)NR⁷R⁸, aryl, aryloxy, arylcarbonyl, C₃₋₇cycloalkyl, C₃₋₇cycloalkyloxy, phthalimide-2-yl, Het³ and C(=O)Het³;

R¹² and R¹³ are each independently selected from hydrogen, C₁₋₄alkyl, hydroxyC₁₋₄alkyl,

5 dihydroxyC₁₋₄alkyl, phenyl, phenylC₁₋₄alkyl, C₁₋₄alkyloxyC₁₋₄alkyl, C₁₋₄alkylcarbonyl, phenylcarbonyl, C₁₋₄alkylcarbonyloxyC₁₋₄alkylcarbonyl, hydroxyC₁₋₄alkylcarbonyl, C₁₋₄alkyloxycarbonylcarbonyl, mono- or di(C₁₋₄alkyl)aminoC₁₋₄alkyl, phenylamino-carbonyl, phenylaminothiocarbonyl, C₃₋₇cycloalkyl, pyridinylC₁₋₄alkyl and R⁶;

10 aryl represents phenyl optionally substituted with one, two or three substituents each independently selected from nitro, azido, halo, hydroxy, C₁₋₄alkyl, C₁₋₄alkyloxy, polyhaloC₁₋₄alkyl, NR⁹R¹⁰, R⁶, phenyl, Het³ and C₁₋₄alkyl substituted with NR⁹R¹⁰;

Het¹ represents a heterocycle selected from pyrrolyl, pyrrolinyl, imidazolyl, imidazoliny, pyrazolyl, pyrazoliny, triazolyl, tetrazolyl, furanyl, tetrahydrofuranyl, thienyl, thiolanyl, dioxolanyl, oxazolyl, oxazoliny, isoxazolyl, thiazolyl, thiazoliny, isothiazolyl, 15 thiadiazolyl, oxadiazolyl, pyridinyl, pyrimidinyl, pyrazinyl, pyranyl, pyridazinyl, pyrrolidinyl, piperidinyl, piperazinyl, morpholinyl, thiomorpholinyl, dioxanyl, dithianyl, trithianyl, triazinyl, benzothienyl, isobenzothienyl, benzofuranyl, isobenzofuranyl, benzothiazolyl, benzoxazolyl, indolyl, isoindolyl, indoliny, purinyl, 1H-pyrazolo[3,4-d]pyrimidinyl, benzimidazolyl, quinolyl, isoquinolyl, cinnoliny, phtalazinyl, 20 quinazoliny, quinoxaliny, thiazolopyridiny, oxazolopyridiny, imidazo[2,1-b]thiazolyl; wherein said heterocycles each independently may optionally be substituted with one, or where possible, two or three substituents each independently selected from Het², R¹¹ and C₁₋₄alkyl optionally substituted with Het² or R¹¹;

Het² represents a monocyclic heterocycle selected from pyrrolyl, pyrroliny, imidazolyl, 25 imidazoliny, pyrazolyl, pyrazoliny, triazolyl, tetrazolyl, furanyl, tetrahydrofuranyl, thienyl, thiolanyl, dioxolanyl, oxazolyl, oxazoliny, isoxazolyl, thiazolyl, thiazoliny, isothiazolyl, thiadiazolyl, oxadiazolyl, pyridinyl, pyrimidinyl, pyrazinyl, pyranyl, pyridazinyl, dioxanyl, dithianyl, trithianyl and triazinyl; wherein said monocyclic heterocycles each independently may optionally be substituted with one, or where 30 possible, two or three substituents each independently selected from R¹¹ and C₁₋₄alkyl optionally substituted with R¹¹;

Het³ represents a monocyclic heterocycle selected from pyrrolidinyl, piperidinyl, piperazinyl, morpholinyl, thiomorpholinyl; wherein said monocyclic heterocycles each 35 independently may optionally be substituted with, where possible, one, two or three substituents each independently selected from C₁₋₄alkyl, C₁₋₄alkyloxy, carboxyl, C₁₋₄alkyloxycarbonyl, C₁₋₄alkylcarbonyl, phenylC₁₋₄alkyl, piperidinyl, NR¹²R¹³, R⁶ and C₁₋₄alkyl substituted with R⁶ or NR¹²R¹³.

As used in the foregoing definitions and hereinafter, halo is generic to fluoro, chloro, bromo and iodo; C₃₋₇cycloalkyl is generic to cyclopropyl, cyclobutyl, cyclopentyl, cyclohexyl and cycloheptyl; C₁₋₄alkyl defines straight and branched chain saturated hydrocarbon radicals having from 1 to 4 carbon atoms such as, for example, methyl, ethyl, propyl, butyl, 1-methylethyl, 2-methylpropyl, 2,2-dimethylethyl and the like; C₁₋₆alkyl is meant to include C₁₋₄alkyl and the higher homologues thereof having 5 or 6 carbon atoms such as, for example, pentyl, 2-methylbutyl, hexyl, 2-methylpentyl and the like; polyhaloC₁₋₄alkyl is defined as polyhalosubstituted C₁₋₄alkyl, in particular C₁₋₄alkyl substituted with 1 to 6 halogen atoms, more in particular difluoro- or trifluoromethyl; polyhaloC₁₋₆alkyl is defined as polyhalosubstituted C₁₋₆alkyl.

Het¹, Het² and Het³ are meant to include all the possible isomeric forms of the heterocycles mentioned in the definition of Het¹, Het² or Het³, for instance, pyrrolyl also includes 2*H*-pyrrolyl; triazolyl includes 1,2,4-triazolyl and 1,3,4-triazolyl; oxadiazolyl includes 1,2,3-oxadiazolyl, 1,2,4-oxadiazolyl, 1,2,5-oxadiazolyl and 1,3,4-oxadiazolyl; thiadiazolyl includes 1,2,3-thiadiazolyl, 1,2,4-thiadiazolyl, 1,2,5-thiadiazolyl and 1,3,4-thiadiazolyl; pyranlyl includes 2*H*-pyranlyl and 4*H*-pyranlyl.

The heterocycles represented by Het¹, Het² and Het³ may be attached to the remainder of the molecule of formula (I) through any ring carbon or heteroatom as appropriate. Thus, for example, when the heterocycle is imidazolyl, it may be a 1-imidazolyl, 2-imidazolyl, 4-imidazolyl and 5-imidazolyl; when it is thiazolyl, it may be 2-thiazolyl, 4-thiazolyl and 5-thiazolyl; when it is triazolyl, it may be 1,2,4-triazol-1-yl, 1,2,4-triazol-3-yl, 1,2,4-triazol-5-yl, 1,3,4-triazol-1-yl and 1,3,4-triazol-2-yl; when it is benzthiazolyl, it may be 2-benzthiazolyl, 4-benzthiazolyl, 5-benzthiazolyl, 6-benzthiazolyl and 7-benzthiazolyl.

The pharmaceutically acceptable addition salts as mentioned hereinabove are meant to comprise the therapeutically active non-toxic acid addition salt forms which the compounds of formula (I) are able to form. The latter can conveniently be obtained by treating the base form with such appropriate acids as inorganic acids, for example, hydrohalic acids, e.g. hydrochloric, hydrobromic and the like; sulfuric acid; nitric acid; phosphoric acid and the like; or organic acids, for example, acetic, propanoic, hydroxyacetic, 2-hydroxypropanoic, 2-oxopropanoic, ethanedioic, propanedioic, butanedioic, (Z)-2-butenedioic, (E)-2-butenedioic, 2-hydroxybutanedioic, 2,3-dihydroxybutanedioic, 2-hydroxy-1,2,3-propanetricarboxylic, methanesulfonic, ethanesulfonic, benzenesulfonic, 4-methylbenzenesulfonic, cyclohexanesulfamic, 2-hydroxybenzoic, 4-amino-2-hydroxybenzoic and the like acids. Conversely the salt form can be converted by

treatment with alkali into the free base form.

The compounds of formula (I) containing acidic protons may be converted into their therapeutically active non-toxic metal or amine addition salt forms by treatment with appropriate organic and inorganic bases. Appropriate base salt forms comprise, for example, the ammonium salts, the alkali and earth alkaline metal salts, e.g. the lithium, sodium, potassium, magnesium, calcium salts and the like, salts with organic bases, e.g. the benzathine, *N*-methyl-D-glucamine, 2-amino-2-(hydroxymethyl)-1,3-propanediol, hydrabamine salts, and salts with amino acids such as, for example, arginine, lysine and the like. Conversely the salt form can be converted by treatment with acid into the free acid form.

The term addition salt also comprises the hydrates and solvent addition forms which the compounds of formula (I) are able to form. Examples of such forms are e.g. hydrates, alcoholates and the like.

The *N*-oxide forms of the present compounds are meant to comprise the compounds of formula (I) wherein one or several nitrogen atoms are oxidized to the so-called *N*-oxide. For example, one or more nitrogen atoms of any of the heterocycles in the definition of Het¹, Het² and Het³ may be *N*-oxidized.

Some of the compounds of formula (I) may also exist in their tautomeric forms. Such forms although not explicitly indicated in the above formula are intended to be included within the scope of the present invention. For example, a hydroxy substituted triazine moiety may also exist as the corresponding triazinone moiety; a hydroxy substituted pyrimidine moiety may also exist as the corresponding pyrimidinone moiety.


The term "stereochemically isomeric forms" as used hereinbefore defines all the possible stereoisomeric forms in which the compounds of formula (I) can exist. Unless otherwise mentioned or indicated, the chemical designation of compounds denotes the mixture of all possible stereochemically isomeric forms, said mixtures containing all diastereomers and enantiomers of the basic molecular structure. More in particular, stereogenic centers may have the R- or S-configuration, used herein in accordance with Chemical Abstracts nomenclature. Stereochemically isomeric forms of the compounds of formula (I) are obviously intended to be embraced within the scope of this invention.

The compounds of formula (I) and some of the intermediates in the present invention contain one or more asymmetric carbon atoms. The pure and mixed stereochemically isomeric forms of the compounds of formula (I) are intended to be embraced within the

scope of the present invention.

Whenever used hereinafter, the term "compounds of formula (I)" is meant to also include their *N*-oxide forms, their pharmaceutically acceptable addition salts, and their stereochemically isomeric forms.

The numbering of the phenyl ring bearing substituent R^4 is given hereinbelow and is used herein as such when indicating the position of the R^4 substituents on said phenyl ring, unless otherwise indicated.



The carbon atom bearing the two phenyl rings and the R^1 and $-X-R^2$ substituents will be referred herein as the central carbon atom.

A special group of compounds are those compounds of formula (I) wherein R^1 represents hydrogen, hydroxy, halo, amino, mono- or di(C_{1-4} alkyl)amino, C_{1-6} alkyl, C_{1-6} alkyloxy, C_{3-7} cycloalkyl, aryl or aryl C_{1-6} alkyl; R^2 represents aryl; Het¹; C_{3-7} cycloalkyl; C_{1-6} alkyl or C_{1-6} alkyl substituted with one or two substituents selected from hydroxy, cyano, amino, mono- or di(C_{1-4} alkyl)amino, C_{1-6} alkyloxy, C_{1-6} alkyloxycarbonyl, C_{3-7} cycloalkyl, aryl and Het¹; and if X is NR^3 , then R^2 may also represent C_{1-4} alkylcarbonyl or arylcarbonyl; each R^4 independently represents halo, polyhalo C_{1-6} alkyl, C_{1-6} alkyl, hydroxy, C_{1-6} alkyloxy, C_{1-6} alkylcarbonyloxy, mercapto, C_{1-6} alkylthio, C_{1-6} alkylsulfonyl, C_{1-6} alkylsulfinyl, polyhalo C_{1-6} alkylsulfonyl, aryl, cyano, nitro, amino, mono- and di(C_{1-6} alkyl)amino or (C_{1-6} alkylcarbonyl)amino; each R^5 independently represents halo, polyhalo C_{1-6} alkyl, C_{1-6} alkyl, hydroxy, C_{1-6} alkyloxy, C_{1-6} alkylcarbonyloxy, mercapto, C_{1-6} alkylthio, C_{1-6} alkylsulfonyl, C_{1-6} alkylsulfinyl, polyhalo C_{1-6} alkylsulfonyl, aryl, cyano, nitro, amino, mono- and di(C_{1-6} alkyl)amino or (C_{1-6} alkylcarbonyl)amino; aryl represents phenyl or phenyl substituted with one, two or three substituents selected from halo, hydroxy, C_{1-4} alkyl, C_{1-4} alkyloxy, polyhalo C_{1-4} alkyl, amino, mono- or di(C_{1-4} alkyl)amino and phenyl; Het¹ represents a heterocycle selected from pyrrolyl, pyrrolinyl, imidazolyl, imidazoliny, pyrazolyl, pyrazolinyl, triazolyl, tetrazolyl, furanyl, tetrahydrofuranyl, thienyl, thiolanyl, dioxolanyl, oxazolyl, oxazoliny, isoxazolyl, thiazolyl, thiazolinyl, isothiazolyl, thiadiazolyl, oxadiazolyl, pyridinyl, pyrimidinyl, pyrazinyl, pyranyl, pyridazinyl, piperidinyl,

piperazinyl, morpholinyl, thiomorpholinyl, dioxanyl, dithianyl, trithianyl, triazinyl, benzothienyl, isobenzothienyl, benzofuranyl, isobenzofuranyl, benzthiazolyl, benzoxazolyl, indolyl, isoindolyl, indolinyl, purinyl, 1*H*-pyrazolo[3,4-*d*]pyrimidinyl, benzimidazolyl, quinolyl, isoquinolyl, cinnolinyl, phtalazinyl, quinazolinyl, quinoxalinyl
 5 and thiazolopyridinyl; said heterocycles each independently may be substituted with one, or where possible, two or three R¹¹ substituents, each R¹¹ independently being selected from hydroxy, mercapto, cyano, nitro, C₁₋₄alkyl, C₁₋₄alkyloxy, amino, mono- or di(C₁₋₄alkyl)amino, mono- or di(C₁₋₄alkyl)aminocarbonyl, mono- or di(aryl)amino, halo, polyhaloC₁₋₄alkyl, C₁₋₄alkyloxycarbonyl, aryl, furanyl, thienyl, pyridinyl,
 10 piperidinyl, C₁₋₄alkyl-carbonylpiperidinyl and C₁₋₄alkyl substituted with C₁₋₄alkyloxy, aryl, hydroxy, piperidinyl, amino, mono- or di(C₁₋₄alkyl)amino or C₃₋₇cycloalkyl.

An interesting group of compounds are those compounds of formula (I) wherein the 6-azauracil moiety is connected to the phenyl ring in the para or meta position relative
 15 to the central carbon atom; preferably in the para position.

Suitably, *p* is 0, 1 or 2; preferably 1 or 2.

Suitably, *q* is 0, 1 or 2; preferably 1 or 2.

Suitably, R¹ represents hydrogen, hydroxy, halo, amino, C₁₋₆alkyl, C₁₋₆alkyloxy or
 20 mono- or di(C₁₋₄alkyl)aminoC₁₋₄alkylamino; in particular, hydrogen, methyl and hydroxy.

Suitably, R² represents aryl, Het¹, C₃₋₇cycloalkyl, C₁₋₆alkyl or C₁₋₆alkyl substituted with one or two substituents selected from hydroxy, cyano, amino, mono- or di(C₁₋₄alkyl)-amino, C₁₋₆alkyloxy, C₁₋₆alkylsulfonyloxy, C₁₋₆alkyloxycarbonyl, aryl, Het¹ and
 25 Het¹thio; and if X is NR³, then R² may also represent arylcarbonyl.

Suitably, R³ represents hydrogen or methyl.

Suitably, each R⁴ independently represents C₁₋₆alkyl, halo, polyhaloC₁₋₆alkyl or C₁₋₆alkyloxy.

Suitably, each R⁵ independently represents C₁₋₆alkyl, halo or C₁₋₆alkyloxy.

30 Suitably, each R⁶ independently represents C₁₋₆alkylsulfonyl, aminosulfonyl or phenylC₁₋₄alkylsulfonyl.

Suitably, each R⁷ and each R⁸ are independently selected from hydrogen, C₁₋₄alkyl, hydroxyC₁₋₄alkyl, dihydroxyC₁₋₄alkyl, aryl, arylC₁₋₄alkyl, C₁₋₄alkyloxyC₁₋₄alkyl, mono- or di(C₁₋₄alkyl)aminoC₁₋₄alkyl, arylaminocarbonyl, arylaminothiocarbonyl, C₃₋₇cycloalkyl,
 35 pyridinylC₁₋₄alkyl, Het³ and R⁶.

Suitably, R⁹ and R¹⁰ are each independently selected from hydrogen, C₁₋₄alkyl, C₁₋₄alkylcarbonyloxyC₁₋₄alkylcarbonyl, hydroxyC₁₋₄alkylcarbonyl, C₁₋₄alkyloxy

carbonylcarbonyl, Het³ aminothiocarbonyl and R⁶.

Suitably, R¹² and R¹³ are each independently selected from hydrogen and C₁₋₄alkyl.

Suitably, Het¹ represents a heterocycle selected from imidazolyl, triazolyl, furanyl, oxazolyl, thiazolyl, thiazolinyl, thiadiazolyl, oxadiazolyl, pyridinyl, pyrimidinyl, pyrazinyl, 5 piperidinyl, piperazinyl, triazinyl, benzothiazolyl, benzoxazolyl, purinyl, 1*H*-pyrazolo-[3,4-*d*]pyrimidinyl, benzimidazolyl, thiazolopyridinyl, oxazolopyridinyl, imidazo-[2,1-*b*]thiazolyl; wherein said heterocycles each independently may optionally be substituted with one, or where possible, two or three substituents each independently selected from Het², R¹¹ and C₁₋₄alkyl optionally substituted with Het² or R¹¹.

10 Suitably, Het² represents furanyl, thienyl or pyridinyl; wherein said monocyclic heterocycles each independently may optionally be substituted with C₁₋₄alkyl.

Suitably, Het³ represents pyrrolidinyl, piperidinyl, piperazinyl, morpholinyl, thiomorpholinyl; wherein said monocyclic heterocycles each independently may optionally be substituted with, where possible, one, two or three substituents each independently 15 selected from C₁₋₄alkyl, C₁₋₄alkyloxy, C₁₋₄alkyloxycarbonyl, C₁₋₄alkylcarbonyl, phenylC₁₋₄alkyl, piperidinyl, NR¹²R¹³ and C₁₋₄alkyl substituted with NR¹²R¹³.

Particular compounds are those compounds of formula (I) wherein R⁴ and R⁵ each independently are halo, polyhaloC₁₋₆alkyl, C₁₋₆alkyl, C₁₋₆alkyloxy or aryl, more in 20 particular, chloro or trifluoromethyl.


Other particular compounds are those compounds of formula (I) wherein R² represents aryl, Het¹, C₃₋₇cycloalkyl or C₁₋₆alkyl substituted with one or two substituents selected from hydroxy, cyano, amino, mono- or di(C₁₋₄alkyl)amino, C₁₋₆alkyloxy, C₁₋₆alkyl- 25 sulfonyloxy, C₁₋₆alkyloxycarbonyl, C₃₋₇cycloalkyl, aryl, aryloxy, arylthio, Het¹, Het¹oxy and Het¹thio; and if X is O, S or NR³, then R² may also represent aminocarbonyl, aminothiocarbonyl, C₁₋₄alkylcarbonyl, C₁₋₄alkylthiocarbonyl, arylcarbonyl or arylthiocarbonyl; more in particular R² is oxadiazolyl, thiazolyl, pyrimidinyl or pyridinyl; wherein said heterocycles each independently may optionally be substituted with one, or 30 where possible, two or three substituents each independently selected from Het², R¹¹ and C₁₋₄alkyl optionally substituted with Het² or R¹¹.

Yet other particular compounds are those compounds of formula (I) wherein X is O, S, NH or a direct bond, more preferably S or a direct bond, most preferably a direct bond. 35

Preferred compounds are those compounds of formula (I) wherein q is 1 or 2 and one R⁴ substituent, preferably chloro, is in the 4 position.

Other preferred compounds are those compounds of formula (I) wherein p is 1 or 2 and the one or two R⁵ substituents, preferably chloro, are in the ortho position relative to the central carbon atom.


- 5 More preferred compounds are those compounds of formula (I) wherein the 6-azauracil moiety is in the para position relative to the central carbon atom; p is 2 and both R⁵ substituent are chloro positioned ortho relative to the central carbon atom; q is 1 and R⁴ is chloro positioned in the 4 position.
- 10 Most preferred compounds include
- 2-[3,5-dichloro-4-[(4-chlorophenyl)(2-pyrimidinylthio)methyl]phenyl]-1,2,4-triazine-3,5(2*H*,4*H*)-dione;
- 2-[3,5-dichloro-4-[(4-chlorophenyl)[2-(4-pyridinyl)-4-thiazolyl]methyl]phenyl]-1,2,4-triazine-3,5(2*H*,4*H*)-dione monohydrochloride.monohydrate;
- 15 2-[3,5-dichloro-4-[(4-chlorophenyl)(5-phenyl-1,3,4-oxadiazol-2-yl)methyl]phenyl]-1,2,4-triazine-3,5(2*H*,4*H*)-dione;
- 2-[3,5-dichloro-4-[(4-chlorophenyl)[4-(2-chlorophenyl)-2-thiazolyl]methyl]phenyl]-1,2,4-triazine-3,5(2*H*,4*H*)-dione;
- 20 2-[3,5-dichloro-4-[(4-chlorophenyl)[4-(3-fluorophenyl)-2-thiazolyl]methyl]phenyl]-1,2,4-triazine-3,5(2*H*,4*H*)-dione;
- 2-[3,5-dichloro-4-[(4-chlorophenyl)(2-pyridinylthio)methyl]phenyl]-1,2,4-triazine-3,5(2*H*,4*H*)-dione; the *N*-oxides, the pharmaceutically acceptable addition salts and the stereochemically isomeric forms thereof.
- 25 In order to simplify the structural representation of the compounds of formula (I), the group




will hereinafter be represented by the symbol D.

- 30 Compounds of formula (I) can generally be prepared by reacting an intermediate of formula (II) wherein W¹ is a suitable leaving group such as, for example, a halogen atom, with an appropriate reagent of formula (III).

-10-




- Said reaction may be performed in a reaction-inert solvent such as, for example, acetonitrile, *N,N*-dimethylformamide, acetic acid, tetrahydrofuran, ethanol or a mixture thereof. Alternatively, in case the reagent of formula (III) acts as a solvent, no additional reaction-inert solvent is required. The reaction is optionally carried out in the presence of a base such as, for example, 1,8-diazabicyclo[5.4.0]undec-7-ene, sodium bicarbonate, sodiummethanolate and the like. Convenient reaction temperatures range between - 70°C and reflux temperature.
- In this and the following preparations, the reaction products may be isolated from the reaction medium and, if necessary, further purified according to methodologies generally known in the art such as, for example, extraction, crystallization, distillation, trituration and chromatography.
- Alternatively, compounds of formula (I) may generally be prepared by cyclizing an intermediate of formula (IV) wherein L is a suitable leaving group such as, for example, C₁₋₆alkyloxy or halo, and E represents an appropriate electron attracting group such as, for example, an ester, an amide, a cyanide, C₁₋₆alkylsulfonyloxy and the like groups; and eliminating the group E of the thus obtained triazinedione of formula (V). Said reaction procedure is analogous to the one described in EP-A-0,170,316.



- Some of the compounds and intermediates of the present invention can be prepared according to or analogous to the procedures described in EP-A-0,170,316 and EP-A-0,232,932.
- For instance, scheme 1 depicts a reaction pathway for the preparation of compounds of formula (I) wherein R¹ is hydrogen and X is a direct bond, said compounds being represented by formula (I-a-1). A ketone of formula (VI) can be reacted with a reagent of formula (VII) wherein W² is a suitable leaving group such as, for example, a halogen,

in a reaction-inert solvent such as, for example, tetrahydrofuran, diethylether, and in the presence of a suitable base such as, for example, butyl lithium, thus forming an intermediate of formula (VIII). The hydroxy group of the intermediates of formula (VIII) may be eliminated by using a suitable reagent such as for example, formamide in acetic acid or triethylsilane in trifluoroacetic acid, thus obtaining an intermediate of formula (IX) of which the nitro group may subsequently be reduced to an amino group which in turn may then be converted to the 6-azauracil group as described in EP-A-0,170,316, thus obtaining compounds of formula (I-a-1).

Scheme 1



10

In addition to the reaction procedure shown in scheme 1, other compounds of formula (I) wherein X is a direct bond may be prepared starting from a ketone of formula (X) (Scheme 2). Reacting said ketone of formula (X) with an intermediate of formula (III) wherein X is a direct bond, said intermediates being represented by formula (III-a), results in a compound of formula (I) wherein R^1 is hydroxy and X is a direct bond, said compounds being represented by formula (I-a-2). Said reaction may be performed in a reaction-inert solvent such as, for example, tetrahydrofuran, diethylether, diisopropylacetamide or a mixture thereof, in the presence of a base such as, for example, butyl lithium, and optionally in the presence of chlorotriethylsilane. Alternatively, intermediate of formula (III-a) may first be transformed into a Grignard reagent, which may then be reacted with the ketone of formula (X). Said compounds of formula (I-a-2) may further be converted to compounds of formula (I) wherein R^1 is a C_{1-6} alkyloxy group represented by formula (I-a-3) using art-known group transformation reactions. The compounds of formula (I-a-2) may also be converted to compounds of formula (I) wherein R^1 is halo, said compounds being represented by formula (I-a-4). A convenient procedure is converting the hydroxy group to a chlorine atom using a suitable reagent

25

such as, for example, thionyl chloride. Said compounds of formula (I-a-4) may further be converted to compounds of formula (I) wherein R^1 is amino, said compounds being represented by formula (I-a-5), using ammonia or a functional derivative thereof, in a reaction-inert solvent such as, for example, tetrahydrofuran; or may be converted to

5 compounds of formula (I-a-3) using art-known group transformation reactions. Reducing the ketone of formula (X) to its corresponding hydroxy derivative of formula (XI) using a suitable reducing agent such as, for example, sodiumborohydride in a reaction-inert solvent such as for example, water, an alcohol, tetrahydrofuran or a mixture thereof; subsequently converting said hydroxy group to a suitable leaving group

10 W^4 being for example a halogen, thus obtaining an intermediate of formula (XII), and finally reacting said intermediate of formula (XII) with an intermediate of formula (III) in a suitable solvent such as, for example, tetrahydrofuran, *N,N*-dimethylformamide, acetonitrile, acetic acid, ethanol or a mixture thereof, and optionally in the presence of a suitable base such as, for example, 1,8-diazabicyclo[5.4.0]undec-7-ene or

15 sodiumbicarbonate, will result in a compound of formula (I) wherein R^1 is hydrogen, said compounds being represented by formula (I-b). Alternatively, intermediates of formula (XI) can be directly transformed to compounds of formula (I-b) wherein X is S, said compounds being represented by formula (I-b-1), using


20 a suitable mercapto containing reagent of formula R^2-SH in a suitable reaction solvent such as, for example, trifluoroacetic acid, methanesulfonic acid, trifluoromethanesulfonic acid or the like.

Also starting from a ketone of formula (X), compounds of formula (I) may be prepared wherein R^1 is hydrogen and $-X-R^2$ is $-NH-C(=O)-$ (aryl or C_{1-6} alkyl), said compounds being represented by formula (I-c). To that effect, a ketone of formula (X) is reacted with

25 formamide in formic acid or a functional derivative thereof, at elevated temperatures. The resulting intermediate of formula (XIII) is hydrolysed to the corresponding amine of formula (XIV), which may then be further reacted with an intermediate of formula (XV) wherein W^3 is a suitable leaving group, in the presence of a suitable base, such as, for example pyridine, optionally in the presence of a reaction-inert solvent such as, for

30 example, dichloromethane.


Scheme 2



Compounds of formula (I) wherein X is a direct bond and R^2 is a heterocycle, said compounds being generally represented by formula (I-d), can conveniently be prepared by cyclization of the appropriate intermediate. Both intramolecular and intermolecular cyclization procedures are feasible and scheme 3 lists several examples.


Starting point is the conversion of the cyano group of an intermediate of formula (XVI) to a carboxyl group thus forming intermediates of formula (XVII) using art-known techniques such as, for example, using a combination of sulfuric- and acetic acid in water, which in turn may be further reacted to acyl halides of formula (XVIII), for instance, the acyl chloride derivative may be prepared using thionyl chloride.

Scheme 3




- The intermediate of formula (XVIII) may be reacted with an intermediate of formula (XIX-a) wherein Y is O, S or NR^3 , to form an intermediate of formula (XX) in the presence of a base such as, for example, pyridine. Said intermediate of formula (XX) may further be cyclized to a compound of formula (I) wherein $-\text{X-R}^2$ is an optionally substituted benzothiazole or benzoxazole, said compounds being represented by formula (I-d-1), in the presence of a suitable solvent such as, for example, acetic acid, at an elevated temperature, preferably at reflux temperature. It may be convenient to prepare compounds of formula (I-d-1) without isolating intermediates of formula (XX).
- Analogously, an intermediate of formula (XVIII) may be reacted with an intermediate of formula (XIX-b) to form an intermediate of formula (XXI) which is cyclized to a compound of formula (I) wherein $-\text{X-R}^2$ is an optionally 3-substituted 1,2,4-oxadiazole, said compounds being represented by formula (I-d-2), in a reaction-inert solvent such as, for example, toluene, at an elevated temperature, preferably at reflux temperature.
- Also analogously, an intermediate of formula (XVIII) may be reacted with an intermediate of formula (XIX-c) wherein Y is O, S or NR^3 , to form an intermediate of formula (XXII) which is cyclized to a compound of formula (I) wherein $-\text{X-R}^2$ is an optionally substituted 1,2,4-triazole, 1,3,4-thiadiazole or 1,3,4-oxadiazole, said compounds being represented by formula (I-d-3), in a suitable solvent such as, for example, phosphorousoxychloride.
- Also analogously, an intermediate of formula (XVIII) may be reacted with an intermediate of formula (XIX-d) wherein Y is O, S or NR^3 , to form an intermediate of formula (XXIII) which is cyclized to a compound of formula (I) wherein $-\text{X-R}^2$ is an optionally amino substituted 1,2,4-triazole, 1,3,4-thiadiazole or 1,3,4-oxadiazole, said compounds being represented by formula (I-d-4) in a reaction-inert solvent such as, for example, toluene, and in the presence of an acid; or, which is cyclized to a compound of formula (I) wherein $-\text{X-R}^2$ is a disubstituted 1,3,4-triazole, said compounds being represented by formula (I-d-5).
- The nitrile derivative of formula (XVI) may also be reacted with hydroxylamine hydrochloride or a functional derivative thereof, thus forming an intermediate of formula (XXIV) which may be reacted with an intermediate of formula (XXV) to form a compound of formula (I) wherein $-\text{X-R}^2$ is an optionally 5-substituted 1,2,4-triazole, 1,2,4-thiadiazole or 1,2,4-oxadiazole, said compounds being represented by formula (I-d-6), in a reaction-inert solvent such as, for example, methanol, butanol or a mixture thereof, and in the presence of a base such as, for example, sodium methanolate.
- Compounds of formula (I-d) wherein the heterocycle is substituted 2-thiazolyl, said compounds being represented by formula (I-d-7), can be prepared by reacting an

- intermediate of formula (XVI) with hydrosulfide or a functional derivative thereof, in a reaction inert solvent such as, for example, pyridine, optionally in the presence of a suitable base such as, for example, triethylamine, thus forming an intermediate of formula (XXVI), which may subsequently be reacted with an intermediate of formula (XXVII) or a functional derivative thereof such as the ketal derivative thereof, in a reaction-inert solvent such as, for example, ethanol, and optionally in the presence of an acid such as, for example, hydrogenchloride.




- Compounds of formula (I-d) wherein the heterocycle is substituted 5-thiazolyl and R^1 is hydrogen, said compounds being represented by formula (I-d-8), can be prepared following the reaction procedure depicted in scheme 4.



- Initially, an intermediate of formula (XXVIII) wherein P is a protective group such as, for example, a C_{1-6} alkylcarbonyl group, is reacted with a thiazole derivative of formula (XXIX) in the presence of a suitable base such as, for example, butyl lithium, in a

- reaction inert solvent such as, for example, tetrahydrofuran, thus forming an intermediate of formula (XXX). It may be convenient to perform said reaction under an inert atmosphere at lower temperature, preferably at about -70°C . The hydroxy group and the protective group P of said intermediates (XXX) may be removed using art-
- 5 known procedures such as, for example, stannous chloride and hydrochloric acid in acetic acid, thus forming an intermediate of formula (XXXI), of which the amino group may further be converted to a 6-azauracil moiety according to the procedure described in EP-A-0,170,316, thus forming a compound of formula (I-d-8).
- 10 Also, compounds of formula (I-d) wherein the heterocycle is 4-thiazolyl, said compounds being represented by formula (I-d-9), can be prepared following the reaction procedure depicted in scheme 5.

Scheme 5



- 15 An intermediate of formula (XVIII) is reacted with a Grignard reagent of formula RCH_2MgBr or a functional derivative thereof to form an intermediate of formula (XXXII), which may be halogenated, preferably brominated, in the α -position using a suitable reagent such as trimethylphenylammonium tribromide in tetrahydrofuran, thus
- 20 forming an intermediate of formula (XXXIII). Said intermediate (XXXIII) may then be reacted with a thioamide of formula (XXXIV) to form a compound of formula (I-d-9), in a reaction-inert solvent such as, for example, ethanol, at an elevated temperature, preferably reflux temperature.
- 25 The compounds of formula (I) can also be converted into each other following art-

known procedures of functional group transformation of which some examples are mentioned hereinabove.

The compounds of formula (I) may also be converted to the corresponding *N*-oxide forms following art-known procedures for converting a trivalent nitrogen into its *N*-oxide form. Said *N*-oxidation reaction may generally be carried out by reacting the starting material of formula (I) with 3-phenyl-2-(phenylsulfonyl)oxaziridine or with an appropriate organic or inorganic peroxide. Appropriate inorganic peroxides comprise, for example, hydrogen peroxide, alkali metal or earth alkaline metal peroxides, e.g. sodium peroxide, potassium peroxide; appropriate organic peroxides may comprise peroxy acids such as, for example, benzenecarboperoxoic acid or halo substituted benzenecarboperoxoic acid, e.g. 3-chlorobenzenecarboperoxoic acid, peroalkanoic acids, e.g. peroacetic acid, alkylhydroperoxides, e.g. *t*-butyl hydroperoxide. Suitable solvents are, for example, water, lower alkanols, e.g. ethanol and the like, hydrocarbons, e.g. toluene, ketones, e.g. 2-butanone, halogenated hydrocarbons, e.g. dichloromethane, and mixtures of such solvents.

Pure stereochemically isomeric forms of the compounds of formula (I) may be obtained by the application of art-known procedures. Diastereomers may be separated by physical methods such as selective crystallization and chromatographic techniques, e.g. counter-current distribution, liquid chromatography and the like.

Some of the compounds of formula (I) and some of the intermediates in the present invention may contain an asymmetric carbon atom. Pure stereochemically isomeric forms of said compounds and said intermediates can be obtained by the application of art-known procedures. For example, diastereoisomers can be separated by physical methods such as selective crystallization or chromatographic techniques, e.g. counter current distribution, liquid chromatography and the like methods. Enantiomers can be obtained from racemic mixtures by first converting said racemic mixtures with suitable resolving agents such as, for example, chiral acids, to mixtures of diastereomeric salts or compounds; then physically separating said mixtures of diastereomeric salts or compounds by, for example, selective crystallization or chromatographic techniques, e.g. liquid chromatography and the like methods; and finally converting said separated diastereomeric salts or compounds into the corresponding enantiomers. Pure stereochemically isomeric forms may also be obtained from the pure stereochemically isomeric forms of the appropriate intermediates and starting materials, provided that the intervening reactions occur stereospecifically.

An alternative manner of separating the enantiomeric forms of the compounds of formula (I) and intermediates involves liquid chromatography, in particular liquid chromatography using a chiral stationary phase.

- 5 Some of the intermediates and starting materials as used in the reaction procedures mentioned hereinabove are known compounds and may be commercially available or may be prepared according to art-known procedures.

10 IL-5, also known as eosinophil differentiating factor (EDF) or eosinophil colony stimulating factor (Eo-CSF), is a major survival and differentiation factor for eosinophils and therefore thought to be a key player in eosinophil infiltration into tissues. There is ample evidence that eosinophil influx is an important pathogenic event in bronchial asthma and allergic diseases such as, cheilitis, irritable bowel disease, eczema, urticaria, vasculitis, vulvitis, winterfeet, atopic dermatitis, pollinosis, allergic rhinitis and allergic
15 conjunctivitis; and other inflammatory diseases, such as eosinophilic syndrome, allergic angitis, eosinophilic fasciitis, eosinophilic pneumonia, PIE syndrome, idiopathic eosinophilia, eosinophilic myalgia, Crohn's disease, ulcerative colitis and the like diseases.

- 20 The present compounds also inhibit the production of other chemokines such as monocyte chemotactic protein-1 and -3 (MCP-1 and MCP-3). MCP-1 is known to attract both T-cells, in which IL-5 production mainly occurs, and monocytes, which are known to act synergetically with eosinophils (Carr et al., 1994, Immunology, 91, 3652-3656). MCP-3 also plays a primary role in allergic inflammation as it is known to
25 mobilize and activate basophil and eosinophil leukocytes (Baggiolini et al., 1994, Immunology Today, 15(3), 127-133).

- The present compounds have no or little effect on the production of other chemokines such as IL-1, IL-2, IL-3, IL-4, IL-6, IL-10, γ -interferon (IFN- γ) and granulocyte-macrophage colony stimulating factor (GM-CSF) indicating that the present IL-5
30 inhibitors do not act as broad-spectrum immunosuppressives.

- The selective chemokine inhibitory effect of the present compounds can be demonstrated by *in vitro* chemokine measurements in human blood of which the test
35 results for IL-5 are presented in the experimental part hereinafter. *In vivo* observations such as the inhibition of eosinophilia in mouse ear, the inhibition of blood eosinophilia in the *Ascaris* mouse model; the reduction of serum IL-5 protein production and splenic IL-5 mRNA expression induced by anti-CD3 antibody in mice and the inhibition of

allergen- or Sephadex-induced pulmonary influx of eosinophils in guinea-pig are indicative for the usefulness of the present compounds in the treatment of eosinophil-dependent inflammatory diseases.

- 5 The present inhibitors of IL-5 production are orally active compounds.

In view of the above pharmacological properties, the compounds of formula (I) can be used as a medicine. In particular, the present compounds can be used in the manufacture of a medicament for treating eosinophil-dependent inflammatory diseases as mentioned
10 hereinabove, more in particular bronchial asthma, atopic dermatitis, allergic rhinitis and allergic conjunctivitis.

In view of the utility of the compounds of formula (I), there is provided a method of treating warm-blooded animals, including humans, suffering from eosinophil-dependent
15 inflammatory diseases, in particular bronchial asthma, atopic dermatitis, allergic rhinitis and allergic conjunctivitis. Said method comprises the systemic or topical administration of an effective amount of a compound of formula (I), a *N*-oxide form, a pharmaceutically acceptable addition salt or a possible stereoisomeric form thereof, to warm-blooded animals, including humans.
20

The present invention also provides compositions for treating eosinophil-dependent inflammatory diseases comprising a therapeutically effective amount of a compound of formula (I) and a pharmaceutically acceptable carrier or diluent.

- 25 To prepare the pharmaceutical compositions of this invention, a therapeutically effective amount of the particular compound, in base form or addition salt form, as the active ingredient is combined in intimate admixture with a pharmaceutically acceptable carrier, which may take a wide variety of forms depending on the form of preparation desired for administration. These pharmaceutical compositions are desirably in unitary dosage form
30 suitable, preferably, for systemic administration such as oral, percutaneous, or parenteral administration; or topical administration such as via inhalation, a nose spray, eye drops or via a cream, gel, shampoo or the like. For example, in preparing the compositions in oral dosage form, any of the usual pharmaceutical media may be employed, such as, for example, water, glycols, oils, alcohols and the like in the case of oral liquid preparations
35 such as suspensions, syrups, elixirs and solutions; or solid carriers such as starches, sugars, kaolin, lubricants, binders, disintegrating agents and the like in the case of powders, pills, capsules and tablets. Because of their ease in administration, tablets and capsules represent the most advantageous oral dosage unit form, in which case solid pharmaceutical carriers are obviously employed. For parenteral compositions, the carrier

will usually comprise sterile water, at least in large part, though other ingredients, for example, to aid solubility, may be included. Injectable solutions, for example, may be prepared in which the carrier comprises saline solution, glucose solution or a mixture of saline and glucose solution. Injectable suspensions may also be prepared in which case
5 appropriate liquid carriers, suspending agents and the like may be employed. In the compositions suitable for percutaneous administration, the carrier optionally comprises a penetration enhancing agent and/or a suitable wettable agent, optionally combined with suitable additives of any nature in minor proportions, which additives do not cause any significant deleterious effects on the skin. Said additives may facilitate the administration
10 to the skin and/or may be helpful for preparing the desired compositions. These compositions may be administered in various ways, e.g., as a transdermal patch, as a spot-on or as an ointment. As appropriate compositions for topical application there may be cited all compositions usually employed for topically administering drugs e.g. creams, gellies, dressings, shampoos, tinctures, pastes, ointments, salves, powders and
15 the like. Application of said compositions may be by aerosol, e.g. with a propellant such as nitrogen, carbon dioxide, a freon, or without a propellant such as a pump spray, drops, lotions, or a semisolid such as a thickened composition which can be applied by a swab. In particular, semisolid compositions such as salves, creams, gellies, ointments and the like will conveniently be used.

20 It is especially advantageous to formulate the aforementioned pharmaceutical compositions in dosage unit form for ease of administration and uniformity of dosage. Dosage unit form as used in the specification and claims herein refers to physically discrete units suitable as unitary dosages, each unit containing a predetermined quantity
25 of active ingredient calculated to produce the desired therapeutic effect in association with the required pharmaceutical carrier. Examples of such dosage unit forms are tablets (including scored or coated tablets), capsules, pills, powder packets, wafers, injectable solutions or suspensions, teaspoonfuls, tablespoonfuls and the like, and segregated multiples thereof.

30 In order to enhance the solubility and/or the stability of the compounds of formula (I) in pharmaceutical compositions, it can be advantageous to employ α -, β - or γ -cyclodextrins or their derivatives. Also co-solvents such as alcohols may improve the solubility and/or the stability of the compounds of formula (I) in pharmaceutical
35 compositions. In the preparation of aqueous compositions, addition salts of the subject compounds are obviously more suitable due to their increased water solubility.

Appropriate cyclodextrins are α -, β -, γ -cyclodextrins or ethers and mixed ethers thereof

wherein one or more of the hydroxy groups of the anhydroglucose units of the cyclodextrin are substituted with C₁₋₆alkyl, particularly methyl, ethyl or isopropyl, e.g. randomly methylated β -CD; hydroxyC₁₋₆alkyl, particularly hydroxyethyl, hydroxypropyl or hydroxybutyl; carboxyC₁₋₆alkyl, particularly carboxymethyl or carboxyethyl; C₁₋₆alkylcarbonyl, particularly acetyl; C₁₋₆alkyloxycarbonylC₁₋₆alkyl or carboxy-C₁₋₆alkyloxyC₁₋₆alkyl, particularly carboxymethoxypropyl or carboxyethoxypropyl; C₁₋₆alkylcarbonyloxyC₁₋₆alkyl, particularly 2-acetyloxypropyl. Especially noteworthy as complexants and/or solubilizers are β -CD, randomly methylated β -CD, 2,6-dimethyl- β -CD, 2-hydroxyethyl- β -CD, 2-hydroxyethyl- γ -CD, 2-hydroxypropyl- γ -CD and (2-carboxymethoxy)propyl- β -CD, and in particular 2-hydroxypropyl- β -CD (2-HP- β -CD).

The term mixed ether denotes cyclodextrin derivatives wherein at least two cyclodextrin hydroxy groups are etherified with different groups such as, for example, hydroxypropyl and hydroxyethyl.

The average molar substitution (M.S.) is used as a measure of the average number of moles of alkoxy units per mole of anhydroglucose. The M.S. value can be determined by various analytical techniques, preferably, as measured by mass spectrometry, the M.S. ranges from 0.125 to 10.

The average substitution degree (D.S.) refers to the average number of substituted hydroxyls per anhydroglucose unit. The D.S. value can be determined by various analytical techniques, preferably, as measured by mass spectrometry, the D.S. ranges from 0.125 to 3.

Due to their high degree of selectivity as IL-5 inhibitors, the compounds of formula (I) as defined above, are also useful to mark or identify receptors. To this purpose, the compounds of the present invention need to be labelled, in particular by replacing, partially or completely, one or more atoms in the molecule by their radioactive isotopes. Examples of interesting labelled compounds are those compounds having at least one halo which is a radioactive isotope of iodine, bromine or fluorine; or those compounds having at least one ¹¹C-atom or tritium atom.

One particular group consists of those compounds of formula (I) wherein R³ and/or R⁴ are a radioactive halogen atom. In principle, any compound of formula (I) containing a halogen atom is prone for radiolabelling by replacing the halogen atom by a suitable isotope. Suitable halogen radioisotopes to this purpose are radioactive iodides, e.g.

^{122}I , ^{123}I , ^{125}I , ^{131}I ; radioactive bromides, e.g. ^{75}Br , ^{76}Br , ^{77}Br and ^{82}Br , and radioactive fluorides, e.g. ^{18}F . The introduction of a radioactive halogen atom can be performed by a suitable exchange reaction or by using any one of the procedures as described hereinabove to prepare halogen derivatives of formula (I).

5

Another interesting form of radiolabelling is by substituting a carbon atom by a ^{11}C -atom or the substitution of a hydrogen atom by a tritium atom.

Hence, said radiolabelled compounds of formula (I) can be used in a process of specifically marking receptor sites in biological material. Said process comprises the steps of (a) radiolabelling a compound of formula (I), (b) administering this radiolabelled compound to biological material and subsequently (c) detecting the emissions from the radiolabelled compound. The term biological material is meant to comprise every kind of material which has a biological origin. More in particular this term refers to tissue samples, plasma or body fluids but also to animals, specially warm-blooded animals, or parts of animals such as organs.

The radiolabelled compounds of formula (I) are also useful as agents for screening whether a test compound has the ability to occupy or bind to a particular receptor site. The degree to which a test compound will displace a compound of formula (I) from such a particular receptor site will show the test compound ability as either an agonist, an antagonist or a mixed agonist/antagonist of said receptor.

When used in *in vivo* assays, the radiolabelled compounds are administered in an appropriate composition to an animal and the location of said radiolabelled compounds is detected using imaging techniques, such as, for instance, Single Photon Emission Computerized Tomography (SPECT) or Positron Emission Tomography (PET) and the like. In this manner the distribution of the particular receptor sites throughout the body can be detected and organs containing said receptor sites can be visualized by the imaging techniques mentioned hereinabove. This process of imaging an organ by administering a radiolabelled compound of formula (I) and detecting the emissions from the radioactive compound also constitutes a part of the present invention.

In general, it is contemplated that a therapeutically effective daily amount would be from 0.01 mg/kg to 50 mg/kg body weight, in particular from 0.05 mg/kg to 10 mg/kg body weight. A method of treatment may also include administering the active ingredient on a regimen of between two or four intakes per day.

Experimental part

Hereinafter, the term 'RT' means room temperature, 'THF' means tetrahydrofuran,

'EtOAc' means ethylacetate, 'DMF' means *N,N*-dimethylformamide, 'MIK' means methylisobutyl ketone, 'DIPE' means diisopropylether, and 'HOAc' means acetic acid.

A. Preparation of the intermediate compounds

Example A.1

- 5 a) A solution of 4-chloro-3-(trifluoromethyl)benzeneacetonitrile (0.114 mol) in THF (100ml) was added dropwise at RT to a solution of 1,2,3-trichloro-5-nitrobenzene (0.114 mol) and *N,N,N*-triethylbenzenemethanaminium chloride (3g) in NaOH (150ml) and THF (100ml). The mixture was stirred for 2 hours, then poured out on ice, acidified with a concentrated HCl solution and extracted with CH₂Cl₂. The organic
10 layer was separated, dried, filtered and the solvent was evaporated. The residue was crystallized from DIPE. The precipitate was filtered off and dried, yielding 40.4 g (86.5%) of (±)-2,6-dichloro-α-[4-chloro-3-(trifluoromethyl)phenyl]-4-nitrobenzene-acetonitrile (interm. 1).
- b) A solution of intermediate (1) (0.0466 mol), iodomethane (0.0606 mol), KOH
15 (0.1864 mol) and *N,N,N*-triethylbenzenemethanaminium chloride (0.0466 mol) in toluene (200ml) was stirred at 50°C for 2 hours. The mixture was poured out into water, acidified with HCl 3N and extracted with CH₂Cl₂. The organic layer was separated, dried, filtered and the solvent was evaporated. The residue was purified by column chromatography over silica gel (eluent: cyclohexane/EtOAc 90/10). The pure
20 fractions were collected and the solvent was evaporated, yielding 11g (55%) of (±)-2,6-dichloro-α-[4-chloro-3-(trifluoromethyl)phenyl]-α-methyl-4-nitrobenzene-acetonitrile (interm. 2).
- c) A mixture of intermediate (2) (0.0259 mol) in methanol (200ml) was hydrogenated at 40°C overnight with platinum-on-charcoal catalyst 1% (1g) as a catalyst in the
25 presence of thiophene 10% in ethanol (1 ml). After uptake of hydrogen (3 equivalents), the catalyst was filtered through celite, washed with CH₃OH and the filtrate was evaporated, yielding 10g (98%) of (±)-4-amino-2,6-dichloro-α-[4-chloro-3-(trifluoromethyl)phenyl]-α-methylbenzeneacetonitrile (interm. 3).

Example A.2

- 30 a) A solution of NaNO₂ (0.0243 mol) in water (10ml) was added dropwise at 5°C to a solution of intermediate (3) (0.0243 mol) in HOAc (75ml) and concentrated HCl (20ml). The mixture was stirred at 0°C for 35 minutes and then added dropwise to a solution of ethyl cyanoacetylcarbamate (0.0326 mol) and sodium acetate (112g) in water (1300ml), cooled to 0°C. The mixture was stirred at 0°C for 45 minutes. The
35 precipitate was filtered off, washed with water and taken up in CH₂Cl₂. The organic layer was separated, washed with water, dried, filtered and the solvent was evaporated,

yielding 15.2g of (±)-ethyl 2-cyano-2-[[3,5-dichloro-4-[1-[4-chloro-3-(trifluoromethyl)phenyl]-1-cyanoethyl]phenyl]hydrozono]-1-oxoethylcarbamate (interm. 4).

- b) A mixture of intermediate (4) (0.0271 mol) and potassiumacetate (0.0285 mol) in HOAc (150ml) was stirred and refluxed for 3 hours and then poured out on ice. The precipitate was filtered off, washed with water and taken up in EtOAc. The organic layer was separated, washed with water, dried, filtered and the solvent was evaporated, yielding 12g (86%) of (±)-2-[3,5-dichloro-4-[1-[4-chloro-3-(trifluoromethyl)phenyl]-1-cyanoethyl]phenyl]-2,3,4,5-tetrahydro-3,5-dioxo-1,2,4-triazine-6-carbonitrile (interm. 5).
- 10 c) A mixture of intermediate (5) (0.0223 mol) in HCl (40ml) and HOAc (150ml) was stirred and refluxed for 3 hours and then poured out into ice water. The precipitate was filtered off, taken up in CH₂Cl₂ and CH₃OH, washed with water, dried, filtered and the solvent was evaporated, yielding 11.4g (96%) of (±)-2-[3,5-dichloro-4-[1-[4-chloro-3-(trifluoromethyl)phenyl]-1-cyanoethyl]phenyl]-2,3,4,5-tetrahydro-3,5-dioxo-1,2,4-
- 15 triazine-6-carboxylic acid (interm. 6).
- d) A mixture of intermediate (6) (0.05 mol) in 2-mercaptoacetic acid (60 ml) was stirred and refluxed for 140 minutes. The reaction mixture was allowed to cool to RT, then poured out into ice-water. The mixture was stirred, then decanted. CH₂Cl₂/CH₃OH (300 ml, 90/10) was added to the residue. The organic layer was separated, washed
- 20 with an aqueous NaHCO₃ solution (200 ml) and with water, then dried, filtered and the solvent was evaporated. The residue was purified over silica gel on a glass filter (eluent: CH₂Cl₂/CH₃OH 99/1). The desired fractions were collected and the solvent was evaporated, yielding 28 g of (±)-2,6-dichloro-α-[4-chloro-3-(trifluoromethyl)phenyl]-4-(4,5-dihydro-3,5-dioxo-1,2,4-triazin-2(3*H*)-yl)-α-methylbenzeneacetonitrile (interm. 7).
- 25 (±)-2-chloro-α-[4-chloro-3-(trifluoromethyl)phenyl]-4-(4,5-dihydro-3,5-dioxo-1,2,4-triazin-2(3*H*)-yl)-5,α-dimethylbenzeneacetonitrile was prepared following the same procedure as described in example A2d (interm. 8).
- e) A mixture of intermediate (7) (0.0106 mol) and triethylamine (0.0106 mol) in pyridine (70 ml) was stirred at 60°C. Gaseous H₂S was bubbled through the mixture for 8 hours.
- 30 The mixture was stirred at 60°C overnight. Gaseous H₂S was bubbled through the mixture for another 10 hours. The mixture was stirred at 60°C overnight. The solvent was evaporated. The residue was taken up in EtOAc, washed with a diluted HCl solution and with water, dried, filtered and the solvent was evaporated. The residue was purified by column chromatography over silica gel (eluent : CH₂Cl₂/CH₃OH 98/2). The
- 35 pure fractions were collected and the solvent was evaporated, yielding 2.5g (45%) of (±)-2,6-dichloro-α-[4-chloro-3-(trifluoromethyl)phenyl]-4-(4,5-dihydro-3,5-dioxo-1,2,4-triazin-2(3*H*)-yl)-α-methylbenzeneethanethioamide (interm. 9).

Following the same procedure there were also prepared :

- (±)-2-chloro- α -[4-chloro-3-(trifluoromethyl)phenyl]-4-[4,5-dihydro-3,5-dioxo-1,2,4-triazin-2(3*H*)-yl]-5, α -dimethylbenzeneethanethioamide (interm. 10);
- (±)-2,6-dichloro- α -(3,4-dichlorophenyl)-4-(4,5-dihydro-3,5-dioxo-1,2,4-triazin-2(3*H*)-yl)- α -methylbenzeneethanethioamide (interm. 11);
- (±)-2-chloro- α -[4-chloro-3-(trifluoromethyl)phenyl]-4-[4,5-dihydro-3,5-dioxo-1,2,4-triazin-2(3*H*)-yl]- α -methylbenzeneethanethioamide (interm. 12);
- (±)-2-chloro- α -(4-chlorophenyl)- α -methyl-4-(4,5-dihydro-3,5-dioxo-1,2,4-triazin-2(3*H*)-yl)benzeneethanethioamide (interm. 13);
- (±)-2,6-dichloro- α -(4-chlorophenyl)-4-(4,5-dihydro-3,5-dioxo-1,2,4-triazin-2(3*H*)-yl)- α -methylbenzeneethanethioamide (interm. 14);
- (±)-2-chloro- α -(4-chlorophenyl)-4-(4,5-dihydro-3,5-dioxo-1,2,4-triazin-2(3*H*)-yl)benzeneethanethioamide (interm. 15).

Example A.3

- a) A mixture of intermediate (1) (0.138 mol) in methanol (300ml) was hydrogenated at RT under a 3 bar pressure for 1 hour with Raney Nickel (50g) as a catalyst in the presence of thiophene solution 10% in ethanol (5ml). After uptake of hydrogen (3 equivalents), the catalyst was filtered through celite, washed with methanol and CH₂Cl₂ and the filtrate was evaporated, yielding 49.5g (94%) of (±)-4-amino-2,6-dichloro- α -[4-chloro-3-(trifluoromethyl)phenyl]benzeneacetonitrile (interm. 16).
- b) (±)-2,6-dichloro- α -[4-chloro-3-(trifluoromethyl)phenyl]-4-(4,5-dihydro-3,5-dioxo-1,2,4-triazin-2(3*H*)-yl)benzeneethanethioamide was prepared following the same procedure as described in A1c and A2a through A2e (interm. 17).
- c) Acetic anhydride (0.1268 mol) was added dropwise at RT to a solution of intermediate (16) (0.0634 mol) in toluene (200ml). The mixture was stirred and refluxed for 3 hours, then cooled, poured out into H₂O and extracted with EtOAc. The organic layer was separated, washed with K₂CO₃ 10% and with H₂O, dried, filtered and the solvent was evaporated, yielding 27.9g (±)-*N*-[3,5-dichloro-4-[[4-chloro-3-(trifluoromethyl)-phenyl]cyanomethyl]phenyl]acetamide (interm. 18; mp. 172°C).

Example A.4

- a) *n*-Butyllithium 1.6 M (0.135 mol) was added dropwise at -70°C under N₂ flow to a solution of 3-bromopyridine (0.11 mol) in 1,1'-oxybisethane (250ml). The mixture was stirred at -70°C for 1 hour. A solution of 2,4'-dichloro-4-nitrodiphenylmethanone (0.0844 mol) in THF (200ml) was added dropwise. The mixture was stirred at -70°C for 3 hours, then poured out into water and extracted with EtOAc. The organic layer was separated, dried, filtered and the solvent was evaporated. The residue was purified

by column chromatography over silica gel (eluent: cyclohexane/EtOAc 60/40 to 100/0). The pure fractions were collected and the solvent was evaporated, yielding 13.7g (43%) of (\pm)- α -(2-chloro-4-nitrophenyl)- α -(4-chlorophenyl)-3-pyridinemethanol (interm. 19).

- b) A mixture of intermediate (19) (0.0373 mol) in methanol (150ml) was hydrogenated
5 at RT under a 3 bar pressure for 4 hours with Raney Nickel (14g) as a catalyst in the presence of thiophene solution 1% in ethanol (2.5ml). After uptake of hydrogen (3 equivalents), the catalyst was filtered through celite and the filtrate was evaporated, yielding 12.06g (94%) of (\pm)- α -(4-amino-2-chlorophenyl)- α -(4-chlorophenyl)-3-pyridinemethanol (interm. 20).
- 10 c) Formamide (60ml) was added to a mixture of intermediate (20) (0.0349 mol) in HOAc (60ml). The mixture was stirred at 150°C for 6 hours, cooled, poured out into ice water, basified with NH_4OH and extracted with CH_2Cl_2 . The organic layer was separated, dried, filtered and the solvent was evaporated, yielding 14.1g of (\pm)-*N*-[3-chloro-4-[(4-chlorophenyl)-3-pyridinylmethyl]phenyl]formamide (interm. 21).
- 15 d) A mixture of intermediate (21) (0.0349 mol) in HCl 6N (150ml) was stirred and refluxed for 4 hours, then cooled, poured out on ice, basified with NH_4OH and extracted with CH_2Cl_2 . The organic layer was separated, dried, filtered and the solvent was evaporated. The residue was purified by column chromatography over silica gel (eluent: $\text{CH}_2\text{Cl}_2/\text{CH}_3\text{OH}/\text{NH}_4\text{OH}$ 98.5/1.5/0.1). The pure fractions were collected and
20 the solvent was evaporated, yielding 7.2g (63%) of (\pm)-3-chloro-4-[(4-chlorophenyl)-3-pyridinylmethyl]benzenamine (interm. 22).
- e) (\pm)-2-[3-chloro-4-[(4-chlorophenyl)-3-pyridinylmethyl]phenyl]-2,3,4,5-tetrahydro-3,5-dioxo-1,2,4-triazine-6-carboxylic acid was prepared following the same procedure as described in A1c and A2a through A2c (interm. 23).

25 Example A.5

- a) A mixture of (\pm)- α -(2-chloro-4-nitrophenyl)- α -(4-chlorophenyl)-1-methyl-1*H*-imidazole-2-methanol (0.0397 mol) and SnCl_2 (0.2382 mol) in HOAc (150ml) and HCl (150ml) was stirred and refluxed for 2 hours, then cooled, poured out on ice, basified with NH_4OH , filtered over celite and extracted with CH_2Cl_2 and CH_3OH . The organic
30 layer was separated, dried, filtered and the solvent was evaporated, yielding 12g (91%) of (\pm)-3-chloro-4-[(4-chlorophenyl)(1-methyl-1*H*-imidazol-2-yl)methyl]benzenamine (interm. 24).
- b) (\pm)-2-[3-chloro-4-[(4-chlorophenyl)(1-methyl-1*H*-imidazol-2-yl)methyl]phenyl]-2,3,4,5-tetrahydro-3,5-dioxo-1,2,4-triazine-6-carboxylic acid (interm. 25);
- 35 (\pm)-2-[3-chloro-4-[(4-chlorophenyl)(1-methyl-1*H*-1,2,4-triazol-5-yl)methyl]phenyl]-2,3,4,5-tetrahydro-3,5-dioxo-1,2,4-triazine-6-carboxylic acid (interm. 26); and

(±)-2-[3-chloro-4-[(4-chlorophenyl)(2-methyl-4-phenyl-5-thiazolyl)methyl]phenyl]-2,3,4,5-tetrahydro-3,5-dioxo-1,2,4-triazine-6-carboxylic acid (interm. 27) were prepared following the same procedure as described in A1c and A2a through A2c.

Example A.6

- 5 a) α-(4-chlorophenyl)-4-pyridinemethanol (0.0512 mol), *N*-(3,5-dichlorophenyl)-acetamide (0.102 mol) and polyphosphoric acid (210g) were stirred at 140°C for 90 minutes. The mixture was cooled to 100°C, poured out into ice water, basified with NH₄OH and extracted with CH₂Cl₂. The organic layer was separated, dried, filtered and the solvent was evaporated. The residue was taken up in 2-propanone and diethyl ether.
- 10 The precipitate was filtered off and the filtrate was evaporated. The residue was purified by column chromatography over silica gel (eluent: CH₂Cl₂/CH₃OH/NH₄OH 97.5/2.5/0.1). The pure fraction was collected and the solvent was evaporated, yielding 17.94g (87%) of (±)-*N*-[3,5-dichloro-4-[(4-chlorophenyl)-4-pyridinylmethyl]phenyl]-acetamide (interm. 28).
- 15 b) The following products were prepared as described in A4c through A4e :
- (±)-2-[3,5-dichloro-4-[(4-chlorophenyl)-4-pyridinylmethyl]phenyl]-2,3,4,5-tetrahydro-3,5-dioxo-1,2,4-triazine-6-carboxylic acid (interm. 29);
- (±)-2-[3,5-dichloro-4-[(4-chlorophenyl)-2-pyridinylmethyl]phenyl]-2,3,4,5-tetrahydro-3,5-dioxo-1,2,4-triazine-6-carboxylic acid (interm. 30);
- 20 (±)-2-[3-chloro-4-[(2-chlorophenyl)-2-pyridinylmethyl]phenyl]-2,3,4,5-tetrahydro-3,5-dioxo-1,2,4-triazine-6-carboxylic acid (interm. 31);
- (±)-2-[3,5-dichloro-4-[(4-chlorophenyl)(1-methyl-1*H*-imidazol-2-yl)methyl]phenyl]-2,3,4,5-tetrahydro-3,5-dioxo-1,2,4-triazine-6-carboxylic acid (interm. 32); and
- (±)-2-[3,5-dichloro-4-[(4-chlorophenyl)-3-pyridinylmethyl]phenyl]-2,3,4,5-tetrahydro-3,5-dioxo-1,2,4-triazine-6-carboxylic acid (interm. 33).
- 25

Example A.7

- a) A mixture of 4-isothiocyanato-2-(trifluoromethyl)-α-[3-(trifluoromethyl)phenyl]-benzeneacetonitrile (0.0516 mol), NaOH solution, 50% (0.155 mol) and *N,N,N*-triethyl-benzenemethanaminium chloride (0.0052 mol) in toluene (250 ml) was stirred for 4
- 30 hours under O₂ at RT. Ice-water and HOAc (9.3 ml) were added. Toluene was added and the reaction mixture was stirred vigorously. The layers were separated. The separated organic layer was dried, filtered and the solvent evaporated. The residue was stirred in hexane. The precipitate was filtered off, washed, and dried, yielding 15.8 g (97.2%) of (4-amino-2-chlorophenyl)[4-chloro-3-(trifluoromethyl)phenyl]methanone
- 35 (interm. 34).

b) (\pm)-2-[3-chloro-4-[4-chloro-3-(trifluoromethyl)benzoyl]phenyl]-1,2,4-triazine-3,5(2*H*,4*H*)-dione (interm. 35) was prepared following the procedures described in A1c and A2a through A2d.

5 c) A mixture of intermediate (35) (0.013 mol) in methanol (50 ml) and THF (50 ml) was stirred at RT. NaBH₄ (0.013 mol) was added portionwise. The reaction mixture was stirred for 1 hour, then acidified (to pH = \pm 6) with concentrated hydrochloric acid. The solvent was evaporated. The residue was purified by column chromatography over silica gel (eluent: CH₂Cl₂/CH₃OH 95/5). The desired fractions were collected and the solvent was evaporated, yielding 5.3 g (94.2%) of (\pm)-2-[3-chloro-4-[[4-chloro-3-(trifluoromethyl)-phenyl]hydroxymethyl]phenyl]-1,2,4-triazine-3,5(2*H*,4*H*)-dione (interm. 36).

10 In a similar way, there was also prepared 2-[3,5-dichloro-4-[(4-fluorophenyl)-hydroxymethyl]phenyl]-1,2,4-triazine-3,5(2*H*,4*H*)-dione (interm. 37).

d) Thionylchloride (5 ml) was added dropwise to a mixture of intermediate (30) (0.012 mol) in CH₂Cl₂ (50 ml), stirred at RT. The resulting reaction mixture was stirred and refluxed for 2 hours. The solvent was evaporated. Toluene was added and azeotroped on the rotary evaporator, yielding 4.9 g (90.4%) of (\pm)-2-[3-chloro-4-[chloro[4-chloro-3-(trifluoromethyl)phenyl]methyl]phenyl]-1,2,4-triazine-3,5(2*H*,4*H*)-dione (interm. 38).

Following the same procedure, there were also prepared :

20 2-[3,5-dichloro-4-[chloro[4-chloro-3-(trifluoromethyl)phenyl]methyl]phenyl]-1,2,4-triazine-3,5(2*H*,4*H*)-dione (interm. 39);

(\pm)-2-[3-chloro-4-[chloro(4-chlorophenyl)-2-thiazolylmethyl]phenyl]-1,2,4-triazine-3,5(2*H*,4*H*)-dione (interm. 40); and

25 (\pm)-2-[4-[(2-benzothiazolyl)chloro(4-chlorophenyl)methyl-3-chlorophenyl]-1,2,4-triazine-3,5(2*H*,4*H*)-dione (interm. 41).

Example A.8

a) K₂CO₃ (0.1786 mol) was added to a solution of intermediate (18) (0.0638 mol) in dimethylsulfoxide (100ml) and water (10ml). Air was bubbled through the mixture for 72 hours. The mixture was poured out into H₂O. The precipitate was filtered off and taken up in EtOAc. The organic solution was washed with H₂O, dried, filtered and the solvent was evaporated. The residue was purified by column chromatography over silica gel (eluent: CH₂Cl₂/CH₃OH 99.25/0.75). The pure fractions were collected and the solvent was evaporated, yielding 18.6g (72%) of *N*-[3,5-dichloro-4-[4-chloro-3-(trifluoromethyl)benzoyl]phenyl]acetamide (interm. 42).

35 b) 2-[3,5-dichloro-4-[4-chloro-3-(trifluoromethyl)benzoyl]phenyl]-2,3,4,5-tetrahydro-3,5-dioxo-1,2,4-triazine-6-carboxylic acid (interm. 43) was prepared following the procedure as described in A6b.

c) 2-[3,5-dichloro-4-[4-chloro-3-(trifluoromethyl)benzoyl]phenyl]-1,2,4-triazine-3,5(2*H*,4*H*)-dione (interm. 44) was prepared following the procedure as described in A2d.

d) 2-[3,5-dichloro-4-[[4-chloro-3-(trifluoromethyl)phenyl]hydroxymethyl]phenyl]-1,2,4-triazine-3,5(2*H*,4*H*)-dione (interm. 45) was prepared following the procedure as described in A7c.

Example A 9

a) A mixture of 4-chloro- α -[2-chloro-4-(4,5-dihydro-3,5-dioxo-1,2,4-triazin-2(3*H*)-yl)-phenyl]- α -methyl-3-(trifluoromethyl)benzeneacetonitrile (0.009 mol) in H₂SO₄ (50ml), HOAc (50ml) and H₂O (40ml) was stirred and refluxed overnight. The mixture was poured out into ice water and extracted with EtOAc. The organic layer was separated, washed with H₂O, dried, filtered and the solvent was evaporated, yielding 4.2g of (\pm)-2-chloro- α -[4-chloro-3-(trifluoromethyl)phenyl]-4-[4,5-dihydro-3,5-dioxo-1,2,4-triazin-2(3*H*)-yl]- α -methylbenzeneacetic acid (interm. 46).

b) A mixture of intermediate (46) (0.009 mol) in thionyl chloride (25ml) was stirred and refluxed for 2.5 hours. The solvent was evaporated, yielding (\pm)-2-chloro- α -[4-chloro-3-(trifluoromethyl)phenyl]-4-[4,5-dihydro-3,5-dioxo-1,2,4-triazin-2(3*H*)-yl]- α -methylbenzeneacetyl chloride (interm. 47).

Following the same procedure, there were also prepared :

(\pm)-2-chloro- α -(4-chlorophenyl)-4-(4,5-dihydro-3,5-dioxo-1,2,4-triazin-2(3*H*)-yl)-benzeneacetyl chloride (interm. 48); and
(\pm)-2-chloro- α -(4-chlorophenyl)-4-(4,5-dihydro-3,5-dioxo-1,2,4-triazin-2(3*H*)-yl)- α -methylbenzeneacetyl chloride (interm. 49).

c) A solution of intermediate (48) (0.011 mol) in 2-propanone (25ml) was added at RT to a solution of *N*-hydroxy benzenecarboximidamide (0.011 mol) and K₂CO₃ (0.011 mol) in 2-propanone (25ml). The mixture was stirred at RT overnight. The precipitate was filtered off, washed with water and taken up in CH₂Cl₂. The organic layer was separated, dried, filtered and the solvent was evaporated, yielding 1.4g (25%) of (\pm)-(iminophenylmethyl)amino 2-chloro-4-(4,5-dihydro-3,5-dioxo-1,2,4-triazin-2(3*H*)-yl)- α -(4-chlorophenyl)benzeneacetate (interm. 50).

Following the same procedure, there was also prepared :
(\pm)-(iminophenylmethyl)amino 2-chloro- α -(4-chlorophenyl)-4-(4,5-dihydro-3,5-dioxo-1,2,4-triazin-2(3*H*)-yl)- α -methylbenzeneacetate (ester) (interm. 51).

d) A solution of intermediate (48) (0.0365 mol) in CH₂Cl₂ (70ml) was added at RT to a solution of 2-aminophenol (0.073 mol) in CH₂Cl₂ (280ml). The mixture was stirred at RT for 12 hours, then washed with HCl 3N and with H₂O, dried, filtered and the solvent was evaporated. The residue was purified by column chromatography over silica gel (eluent : CH₂Cl₂/CH₃OH 98/2). The pure fractions were collected and the solvent was

evaporated, yielding 3.8g (21%) of (\pm) - α -(4-chlorophenyl)-3-chloro-4-(4,5-dihydro-3,5-dioxo-1,2,4-triazin-2(3H)-yl)-N-(2-hydroxyphenyl)benzeneacetamide (interm. 52).

In a similar manner there were also prepared :

- 5 (\pm) -2-chloro- α -(4-chlorophenyl)-4-(4,5-dihydro-3,5-dioxo-1,2,4-triazin-2(3H)-yl)benzeneacetic acid 2-benzoylhydrazide (interm. 53);
- (\pm) -(benzoylamino)-2,6-dichloro- α -(4-chlorophenyl)-4-(4,5-dihydro-3,5-dioxo-1,2,4-triazine-2(3H)-yl)benzeneacetamide (interm. 54);
- (\pm) -2-chloro- α -[4-chloro-3-(trifluoromethyl)phenyl]-4-(4,5-dihydro-3,5-dioxo-1,2,4-triazin-2(3H)-yl)- α -methylbenzeneacetic acid 2-benzoylhydrazide (interm. 55);
- 10 (\pm) -2-chloro- α -(4-chlorophenyl)-4-(4,5-dihydro-3,5-dioxo-1,2,4-triazin-2(3H)-yl)-N-(2-hydroxyphenyl)- α -methylbenzeneacetonitrile (interm. 56);
- (\pm) -2-chloro- α -(4-chlorophenyl)-4-(4,5-dihydro-3,5-dioxo-1,2,4-triazin-2(3H)-yl)-benzeneacetic acid 2-acetylhydrazide (interm. 57);
- (\pm) -2-chloro- α -(4-chlorophenyl)-4-(4,5-dihydro-3,5-dioxo-1,2,4-triazin-2(3H)-yl)-N-(2-phenyl-2-oxoethyl)benzeneacetamide (interm. 58);
- 15 (\pm) -2-[[2-chloro-4-(4,5-dihydro-3,5-dioxo-1,2,4-triazin-3(2H)-yl)phenyl] (4-chlorophenyl)acetyl]-N-phenylhydrazinecarbothioamide (interm. 59);
- (\pm) -2-chloro- α -(4-chlorophenyl)-4-(4,5-dihydro-3,5-dioxo-1,2,4-triazin-2(3H)-yl)- α -methylbenzeneacetic acid 2-benzoylhydrazide (interm. 60); and
- 20 (\pm) -2,6-dichloro- α -(4-chlorophenyl)-4-(4,5-dihydro-3,5-dioxo-1,2,4-triazin-2(3H)-yl)-N-(2-phenyl-2-oxoethyl)benzeneacetamide (interm. 61).

Example A 10

- a) A mixture of 2-[3-chloro-4-[chloro(4-chlorophenyl)methyl]phenyl]-1,2,4-triazine-3,5-(2H,4H)-dione (0.03 mol), thiourea (0.03 mol) and NaHCO₃ (0.03 mol) in DMF
25 (75 ml) was stirred for 18 hours at RT. The solvent was evaporated. The residue was stirred in water, filtered off, washed with water, yielding 12.3 g. (\pm) -[2-chloro-4-(4,5-dihydro-3,5-dioxo-1,2,4-triazin-2(3H)-yl)phenyl]-4-(chloro-phenyl)methyl carbamimidothioate (interm. 62).
- b) A mixture of NaOH (0.25 mol) in water (100 ml) was stirred at RT. (0.03 mol) was
30 added and the resulting reaction mixture was stirred for 18 hours at RT, neutralized, and the precipitate was filtered off and dissolved in CH₂Cl₂. The aqueous phase was separated. The separated organic layer was dried, filtered, and the solvent evaporated. The residue was purified over silica gel on a glass filter (eluent: CH₂Cl₂/CH₃OH/THF 92/3/5). The desired fractions were collected and the solvent was evaporated. The
35 residue was purified by column chromatography over silica gel (eluent: CH₂Cl₂/CH₃OH/THF 92/3/5). The pure fractions were collected and the solvent was evaporated. The residue was stirred in DIPE, filtered off and dried, yielding 4.2 g

(37%) (\pm)-2-[3-chloro-4-[(4-chlorophenyl)mercaptomethyl]phenyl]-1,2,4-triazine-3,5(2*H*,4*H*)-dione (interm. 63).

Example A.11

- 5 a) A mixture of 2-[3-chloro-4-(4-chlorobenzoyl)phenyl]-1,2,4-triazine-3,5(2*H*,4*H*)-dione (0.081 mol) in formic acid (120 ml) and formamide (300 ml) was stirred for 16 hours at 160 °C. The reaction mixture was cooled, poured out into water (600 ml) and the resulting precipitate was filtered off and dried. This fraction was purified by column chromatography over silica gel (eluent: CH₂Cl₂/CH₃OH 95/5). The pure fractions were collected and the solvent was evaporated. The residue was stirred in DIPE, filtered off,
10 and dried, yielding 8.78 g (22.5%) of (\pm)-*N*-[[2-chloro-4-(4,5-dihydro-3,5-dioxo-1,2,4-triazin-2(3*H*)-yl)phenyl](4-chlorophenyl)methyl]formamide (interm. 64).
- b) A mixture of intermediate (64) (0.277 mol) in HCl (200 ml, 36%) and HOAc (1000 ml) was stirred and refluxed for 1 hour. The solvent was evaporated. The residue was taken up into water, then basified with K₂CO₃. The precipitate was filtered off, dried
15 and stirred in boiling ethanol, cooled, filtered off and dried. The precipitate was purified by column chromatography over silica gel (eluent: CH₂Cl₂/CH₃OH 95/5). The pure fractions were collected and the solvent was evaporated. The residue was stirred in boiling CH₃CN, then cooled, filtered off and dried, yielding 1.1 g (\pm)-2-[4-[amino(4-chlorophenyl)methyl]-3-chlorophenyl]-1,2,4-triazine-3,5(2*H*,4*H*)-dione (interm. 65).

20 Example A 12

- NaOCH₃ (0.189 mol; 30 % in CH₃OH) was added to a solution of hydroxylamine (0.189 mol) in ethanol (105ml) The mixture was stirred at RT for 15 minutes and then filtered. The filtrate was added to a mixture of 2-chloro- α -(4-chlorophenyl)-4-(4,5-dihydro-3,5-dioxo-1,2,4-triazin-2(3*H*)-yl)benzeneacetonitrile (0.054 mol) in ethanol
25 (55ml). The mixture was stirred at 60°C for 1 hour, stirred and refluxed for 2 hours and stirred at RT overnight. The solvent was evaporated. The residue was taken up in water and extracted with EtOAc. The organic layer was separated, dried, filtered and the solvent was evaporated, yielding 20.3g of (\pm)-2-chloro- α -(4-chlorophenyl)-4-(4,5-dihydro-3,5-dioxo-1,2,4-triazin-2(3*H*)-yl)-*N*'-hydroxybenzeneethanimidamide
30 (interm. 66).

Example A13

- a) Trifluoro acetic acid (100ml), previously cooled to 5°C, was added dropwise at 0°C/5°C under N₂ flow to (\pm)-1,1-dimethylethyl-2-[2-[2,6-dichloro-4-(4,5-dihydro-3,5-dioxo-1,2,4-triazin-2(3*H*)-yl)phenyl]-2-(4-chlorophenyl)acetyl]hydrazinecarboxylate
35 (0.035 mol). The mixture was allowed to warm to RT and then stirred for 1 hour. The solvent was evaporated. The residue was taken up in H₂O. The precipitate was filtered

-33-

- off, dried, washed with DIPE and dried, yielding 11 g (70%) of R142321 (interm. 67).
- b) A mixture of 3-hydroxy-benzoyl chloride (0.0124 mol) in THF (25ml) was added dropwise at 10°C under N₂ flow to a solution of intermediate 67 (0.0113 mol) and triethylamine (0.0452 mol) in THF (30ml). The mixture was brought to RT. HCl 3N
- 5 was added and the mixture was extracted with CH₂Cl₂. The organic layer was separated, dried, filtered and the solvent was evaporated. The residue was taken up in ethanol. The mixture was filtered and the filtrate was evaporated. The residue was purified by column chromatography over silica gel (eluent: CH₂Cl₂/CH₃OH 95/5). The pure fractions were collected and the solvent was evaporated, yielding 3 g (47%) of (±)-2-[3,5-dichloro-4-
- 10 [1-(4-chlorophenyl)-2-[(3-hydroxybenzoyl)hydrazino]-2-oxoethyl]phenyl]-1,2,4-triazine-3,5(2*H*,4*H*)-dione (interm. 68)

Example A14

- a) A mixture of 2,6-dichloro- α -(4-chlorophenyl)-4-(4,5-dihydro-3,5-dioxo-1,2,4,-triazin-2(3*H*)-yl)-benzeneacetyl chloride (0.05 mol) in THF (200 ml) was stirred at -
- 15 75°C. A solution of chloroethyl magnesium (0.1 mol; 2 M/THF) in THF (50 ml) was added dropwise at -75°C. The reaction mixture was stirred for 90 minutes, then the temperature was raised to -20 °C. A saturated aqueous NH₄Cl solution was added dropwise. Water was added and the product was extracted with CH₂Cl₂. The organic layer was separated, dried, filtered and the solvent was evaporated. The residue was
- 20 filtered over silica gel (eluent: CH₂Cl₂/CH₃OH 98/2). Two fractions were collected and the solvent was evaporated, yielding 3.8 g (±)-2-[3,5-dichloro-4-[1-(4-chlorophenyl)-2-oxobutyl]phenyl]-1,2,4-triazine-3,5(2*H*,4*H*)-dione (interm. 69).
- b) A mixture of intermediate 69 (0.005 mol) in 1,4-dioxane (10 ml) and diethyl ether (20 ml) was stirred at RT. Br₂ (0.005 mol) was added dropwise at RT and the resulting
- 25 reaction mixture was stirred for 15 hours at RT. This mixture was washed 3 times with water and CH₂Cl₂ was added. The separated organic layer was dried, filtered and the solvent evaporated. The residue was dried, yielding 2.6 g (±)-2-[4-[3-bromo-1-(4-chlorophenyl)-2-oxobutyl]-3,5-dichlorophenyl]-1,2,4-triazine-3,5(2*H*,4*H*)-dione (interm. 70).

Example A15

- a) *n*-Butyl lithium (0.045 mol) was added at -70°C under N₂ flow to a solution of 4-phenyl-thiazole (0.045 mol) in diethyl ether (50ml). The mixture was stirred at -70°C for 90 minutes. A solution of 2-[3-chloro-4-(4-chlorobenzoyl)phenyl]-1,2,4-triazine-3,5-(2*H*,4*H*)-dione (0.015 mol) in THF (10ml) was added at -70°C. The mixture was
- 35 stirred at -70°C for 1 hour, then poured out into ice water, neutralized with HCl 3N and extracted with EtOAc. The organic layer was separated, dried, filtered and the solvent

- wasevaporated. The residue was purified by column chromatography oversilica gel (eluent: $\text{CH}_2\text{Cl}_2/\text{CH}_3\text{OH}$ 98/2). The desired fraction was repurified by HPLC (eluent: $\text{CH}_3\text{OH}/(\text{NH}_4\text{OAc}$ 1% in H_2O) 80/20). The pure fractions were collected and the solvent was evaporated, yielding 0.83g (11%) of (\pm)-2-[3-chloro-4-[(4-chlorophenyl)-hydroxy(4-phenyl-2-thiazolyl)methyl]phenyl]-1,2,4-triazine-3,5(2*H*,4*H*)-dione (interm. 71).
- b) A mixture of intermediate 71 (0.0076 mol) in thionyl chloride (35ml) was stirred at 50°C for 4 hours and then brought to RT. The solvent was evaporated, yielding (\pm)-2-[3-chloro-4-[chloro(4-chlorophenyl)(5-chloro-4-phenyl-2-thiazolyl)methyl]phenyl]-1,2,4-triazine-3,5(2*H*,4*H*)-dione (interm. 72).

Example A16

- a) 1-Chloromethoxy-2-methoxy-ethane (0.147 mol) was added dropwise at 15°C to a solution of 3-(3-methoxyphenyl)-8-methyl-8-azabicyclo[3.2.1]octan-3-ol (0.134 mol) and K_2CO_3 (0.134 mol) in DMF (200ml). The mixture was stirred at RT for 24 hours, then poured out into H_2O and extracted with diethyl ether. The organic layer was separated, washed with H_2O , dried, filtered and the solvent was evaporated, yielding 67.27g (\pm)-2-chloro- α -(4-chlorophenyl)-4-[4,5-dihydro-4-[(2-methoxyethoxy)methyl]-3,5-dioxo-1,2,4-triazin-2(3*H*)-yl]benzeneacetonitrile (interm. 73).
- b) NaH (0.063 mol) was added at 10°C under N_2 flow to a solution of intermediate 73 (0.0485 mol) in DMF (100ml). The mixture was stirred for 30 minutes. A solution of 2-chloromethyl-4-phenyl-thiazole (0.063 mol) in DMF (100ml) was added. The mixture was allowed to warm to 15°C over a 2-hour period while stirring, then poured out into ice water and extracted with diethyl ether. The organic layer was separated, washed with H_2O , dried, filtered and the solvent was evaporated. The residue was purified by column chromatography over silica gel (eluent: cyclohexane/EtOAc 65/35). The pure fractions were collected and the solvent was evaporated, yielding 15g (52%) of (\pm)- α -[2-chloro-4-[4,5-dihydro-4-[(2-methoxyethoxy)methyl]-3,5-dioxo-1,2,4-triazin-2(3*H*)-yl]phenyl]- α -(4-chlorophenyl)-4-phenyl-2-thiazolpropanenitrile (interm. 74)
- c) A mixture of intermediate 74 (0.0186 mol) in H_2SO_4 (160ml), acetic acid (160ml) and H_2O (25ml) was stirred and heated for 48 hours. The mixture was cooled and poured out into H_2O . The precipitate was filtered off, taken up in EtOAc and the mixture was separated into its layers. The organic layer was dried, filtered and the solvent was evaporated, to give residue 1. The aqueous layer was evaporated partially and then cooled. The precipitate was filtered off and taken up in EtOAc. The organic solution was dried, filtered and the solvent was evaporated, to give residue 2. Residue 1 and 2 were combined, yielding 8.97g (86%) of (\pm)- α -[2-chloro-4-[4,5-dihydro-3,5-dioxo-

1,2,4-triazin-2(3*H*)-yl]phenyl]- α -(4-chlorophenyl)-4-phenyl-2-thiazolpropanoic acid (interm. 75).

Example A17

- a) NaH (0.0772 mol) was added portionwise at 0°C under N₂ flow to a mixture of 4-chloro-benzeneacetonitrile (0.0643 mol) in DMF (50ml). The mixture was stirred at 0°C under N₂ flow for 1 hour. A mixture of 1,3-dibromo-2-methoxy-5-nitro-benzene (0.0643 mol) in DMF (50ml) was added at 0°C under N₂ flow. The mixture was stirred at RT for 3 hours, hydrolyzed with H₂O and HCl 3N and extracted with EtOAc. The organic layer was separated, dried, filtered and the solvent was evaporated. The residue was purified by column chromatography over silica gel (eluent: CH₂Cl₂/cyclohexane 60/40). The pure fractions were collected and the solvent was evaporated, yielding 12.8g (46%) of (\pm)-2,6-dibromo- α -(4-chlorophenyl)-4-nitrobenzeneacetonitrile (interm. 76).
- b) TiCl₃ (0.13 mol; 15 % in H₂O) was added dropwise at RT to a solution of intermediate 76 (0.026 mol) in THF (200ml). The mixture was stirred at RT for 2 hours, poured out into H₂O and extracted with CH₂Cl₂. The organic layer was separated, washed with H₂O and with K₂CO₃ 10%, dried, filtered and the solvent was evaporated. 2 g of this fraction was crystallized from diethyl ether. The precipitate was filtered off and dried, yielding 1.3g (\pm)-4-amino-2,6-dibromo- α -(4-chlorophenyl)-benzeneacetonitrile (interm. 77).

20 B. Preparation of the final compounds

Example B1

- A mixture of 2-[3-chloro-4-[chloro(4-chlorophenyl)methyl]phenyl]-1,2,4-triazine-3,5(2*H*,4*H*)-dione (0.0075 mol) and 1,8-diazabicyclo[5.4.0]undec-7-ene (0.025 mol) in 2-methylpropanol (25 ml) was stirred for 72 hours at 80 °C. The solvent was evaporated. The residue was purified by column chromatography over silica gel (eluent: CH₂Cl₂/CH₃OH 97.5/2.5). The pure fractions were collected and the solvent was evaporated. The residue was dried, yielding 0.8 g (25%) of (\pm)-2-[3-chloro-4-[(4-chloro-phenyl)(2-methylpropoxy)methyl]phenyl]-1,2,4-triazine-3,5(2*H*,4*H*)-dione (compound 133).

30 Example B2

- a) A mixture of 2-[3-chloro-4-[chloro(4-chlorophenyl)methyl]phenyl]-1,2,4-triazine-3,5(2*H*,4*H*)-dione (0.015 mol) and 2-mercaptopyridine (0.04 mol) in THF (100 ml) was stirred overnight at RT. 1,8-diazabicyclo[5.4.0]undec-7-ene (0.03 mol) was added and the resulting reaction mixture was stirred for 3 hours. NaOH (1 N; 50 ml) was added.
- 35 The mixture was stirred for 5 minutes, then extracted with EtOAc. The separated organic layer was washed with water, dried, filtered and the solvent evaporated. The

- aqueous layers were combined, then acidified (pH = 6) with HCl (1 N). This mixture was extracted with CH₂Cl₂. The separated organic layer was dried, filtered, and the solvent was evaporated. The residue was purified by column chromatography over silica gel (eluent: CH₂Cl₂/THF/CH₃OH 94/5/1). The pure fractions were collected and the solvent was evaporated. The residue was stirred overnight in diethyl ether. The solvent was evaporated. The residue was dried, yielding 2.98 g (43%) (±)-2-[3-chloro-4-[(4-chlorophenyl)(2-pyridinyl-thio)methyl] phenyl]-1,2,4-triazine-3,5(2*H*,4*H*)-dione (compound 93).
- 5 b) (±)-2-[3-chloro-4-[(4-chlorophenyl)(1*H*-imidazol-2-ylthio)methyl]phenyl]-1,2,4-triazine-3,5(2*H*,4*H*)-dione was prepared using the same procedure as in example B2a, but using NaHCO₃ as a base and DMF as a solvent (compound 94).
- 10 c) Sodium (0.075 mol) was added portionwise to ethanol (50 ml) under N₂ atmosphere and this mixture was stirred until complete dissolution. Ethyl 2-amino-3-mercaptopropanoate (0.075 mol) was added and the mixture was stirred for 2 hours at RT. The solvent was evaporated, THF (50 ml) was added to the residue, and a solution of 2-[3-chloro-4-[chloro(4-chlorophenyl)methyl]phenyl]-1,2,4-triazine-3,5(2*H*,4*H*)-dione (0.015 mol) in THF (50 ml) was added. 1,8-diazabicyclo[5.4.0]undec-7-ene (0.03 mol) was added and the resulting reaction mixture was stirred overnight at RT. The solvent was evaporated. The residue was stirred in water and extracted with
- 20 CH₂Cl₂. The organic layer was separated, washed with water, dried, filtered and the solvent was evaporated. This fraction was purified by HPLC over silica gel (eluent: CH₂Cl₂/CH₂OH 97/3). The pure fractions were collected and the solvent was evaporated. The residue was stirred in DIPE, filtered off, washed and dried, yielding (±)-ethyl α-[[[(4-chlorophenyl)[2-chloro-4-(4,5-dihydro-3,5-dioxo-1,2,4-triazin-2(3*H*)-yl)phenyl]methyl]thio]methyl]glycine (compound 95).
- 25 d) A mixture of intermediate 39 (0.00618 mol), 5-amino-4-phenyl-2(3*H*)-thiazolethione (0.00742 mol) and 2,3,4,6,7,8,9,10-octahydropyrimido[1,2-*a*]azepine (0.0124 mol) in dry THF (50 ml) and DMF (50 ml) was stirred and refluxed for four days under N₂ atmosphere. The solvent was evaporated. The residue was taken up into
- 30 CH₂Cl₂/CH₃OH (95/5). The organic solution was washed twice with a saturated aqueous NaCl solution, dried, filtered and the solvent was evaporated. The residue was purified by flash column chromatography over silica gel (eluent: CH₂Cl₂/CH₃OH 97.5/2.5). The desired fractions were collected and the solvent was evaporated. The residue was repurified by HPLC over silica gel (eluent: CH₂Cl₂/CH₃OH 100/0 first 30
- 35 minutes, then 95/5). The pure fractions were collected and the solvent was evaporated. The residue was stirred in boiling CH₃CN, then allowed to cool to RT. The precipitate was filtered off, washed with CH₃CN, then dried, yielding 0.24 g of (±)-2-[3,5-dichloro-4-

[[4-chloro-3-(trifluoromethyl)phenyl][(2,3-dihydro-5-phenyl-2-thioxo-1*H*-imidazol-4-yl)thio]-methyl]phenyl]-1,2,4-triazine-3,5(2*H*,4*H*)-dione (comp. 400).

Example B3

- a) A mixture of 2-[3-chloro-4-[chloro(4-chlorophenyl)methyl]phenyl]-1,2,4-triazine-3,5(2*H*,4*H*)-dione (0.015 mol) and 1-methylpiperazine (0.04 mol) in DMF (100 ml) was stirred for 24 hours at 80 °C. The solvent was evaporated. MIK was added and azeotroped on the rotary evaporator. The residue was stirred in water, then extracted with CH₂Cl₂. The separated organic layer was washed with water, dried, filtered and the solvent evaporated. The residue was purified by column chromatography over silica gel (eluent: CH₂Cl₂/CH₃OH/THF 90/5/5 and CH₂Cl₂/CH₃OH 90/10). The pure fractions were collected and the solvent was evaporated. The residue was stirred overnight in DIPE, then the solvent was evaporated. The residue was crystallized from EtOAc. The precipitate was filtered off, washed with EtOAc, DIPE, then dried, yielding 1.19 g (±)-2-[3-chloro-4-[(4-chlorophenyl)(4-methyl-1-piperazinyl) methyl]phenyl]-1,2,4-triazine-3,5(2*H*,4*H*)-dione (compound 118).
- b) A mixture of 2-[3-chloro-4-[chloro(4-chlorophenyl)methyl]phenyl]-1,2,4-triazine-3,5(2*H*,4*H*)-dione (0.015 mol), 4-hydroxypiperidine (0.02 mol) and sodiumbicarbonate (0.02 mol) in DMF (100ml) was stirred for 16 hours at 80 °C. The mixture was cooled. The solvent was evaporated. The residue was purified by column chromatography over silica gel (eluent: CH₂Cl₂/CH₃OH 95/5). The desired fractions were collected and the solvent was evaporated. The residue was stirred in DIPE, filtered off and dried, yielding 0.070 g (±)-2-[3-chloro-4-[(4-chlorophenyl)(4-hydroxy-1-piperidinyl) methyl]phenyl]-1,2,4-triazine-3,5(2*H*,4*H*)-dione (compound 119).
- c) (±)-2-[3-chloro-4-[(4-chlorophenyl)(2-hydroxyethyl)amino]methyl]phenyl]-1,2,4-triazine-3,5(2*H*,4*H*)-dione was prepared according to the procedure described in example B3a but using CH₃CN as a solvent instead of DMF (compound 51).
- d) Methanol (100 ml) was stirred at RT and sodium (0.09 mol) was added. The mixture was stirred until complete dissolution. (1*H*-imidazol-2-yl)methanamine (0.045 mol) was added. The mixture was stirred for 30 minutes. NaCl was removed by filtration and the filtrate was evaporated. Toluene was added and azeotroped on the rotary evaporator. 2-[3-chloro-4-[chloro(4-chlorophenyl)methyl]phenyl]-1,2,4-triazine-3,5(2*H*,4*H*)-dione (0.015 mol) and acetonitrile (50 ml) were added. The resulting reaction mixture was stirred and refluxed for 20 hours. The solvent was evaporated, the residue was stirred in water, and extracted with CH₂Cl₂/CH₃OH (90/10). The separated organic layer was dried, filtered, and the solvent evaporated. The residue was purified over silica gel on a glass filter (eluent: CH₂Cl₂/CH₃OH 90/10). The pure fractions were collected and the solvent was evaporated. The residue was stirred in CH₃CN, filtered

- off, washed with DIPE, then dried, yielding 1.1 g (16.5%) of (\pm)-2-[3-chloro-4-[(4-chlorophenyl)[(1*H*-imidazol-2-ylmethyl)amino]methyl]phenyl]-1,2,4-triazine-3,5(2*H*,4*H*)-dione (compound 50).
- e) (\pm)-2-[3-chloro-4-[(4-chlorophenyl)(2-pyrimidinylamino)methyl]phenyl]-1,2,4-triazine-3,5(2*H*,4*H*)-dione was prepared according to the procedure described in example B3a but using acetic acid as a solvent instead of DMF (compound 49).
- f) (\pm)-2-[3-chloro-4-[(4-chlorophenyl)[(1-methyl)-4-piperidinyl]amino]methyl]phenyl]-1,2,4-triazine-3,5(2*H*,4*H*)-dione was prepared according to the procedure described in example B3a but using THF as a solvent instead of DMF (compound 48).
- g) A mixture of 2-[4-[chloro(4-chlorophenyl)methyl]-3,5-dichlorophenyl]-1,2,4-triazine-3,5(2*H*,4*H*)-dione (0.00719 mol) and 2-pyrimidinamine (0.00863 mol) was heated for 2 hours at 150 °C in an autoclave. The mixture was cooled to RT. This fraction was taken up into CH₂Cl₂, washed with water, dried, filtered and the solvent was evaporated. The residue was purified by HPLC (eluent: (0.5% NH₄OAc in H₂O)/CH₃OH/CH₃CN gradient elution from 70/15/15 over 0/50/50 to 0/0/100). The desired fractions were collected and the solvent was evaporated. The residue was coevaporated with EtOAc. The residue was stirred in DIPE, filtered off, washed and dried, yielding 0.21 g of (\pm)-2-[3,5-dichloro-4-[(4-chlorophenyl)[(2-pyrimidinyl)amino]methyl]phenyl]-1,2,4-triazine-3,5(2*H*,4*H*)-dione (comp. 413).
- Example B4**
- a) n-Butyllithium, 1.6M (0.0414 mol) was added dropwise at -70°C under N₂ flow to a solution of 1-methyl-1*H*-imidazole (0.0414 mol) in diethyl ether (50ml). The mixture was stirred at -70°C for 90 minutes. A solution of 2-[3-chloro-4-(4-chlorobenzoyl)phenyl]-1,2,4-triazine-3,5(2*H*,4*H*)-dione (0.0138 mol) in THF (100ml) was added dropwise. The mixture was allowed to warm to -40°C, then poured out into ice water, neutralized with HCl 3N and extracted with EtOAc. The organic layer was separated, dried, filtered and the solvent was evaporated. The residue (5.88g) was purified by column chromatography over silica gel (eluent: CH₂Cl₂/CH₃OH 97/3). The pure fractions were collected and the solvent was evaporated. The residue was taken up in CH₃CN and diethyl ether. The precipitate was filtered off and dried, yielding 1.36g (\pm)-2-[3-chloro-4-[(4-chlorophenyl)hydroxy(1-methyl-1*H*-imidazol-2-yl)methyl]phenyl]-1,2,4-triazine-3,5(2*H*,4*H*)-dione monohydrate (compound 120)..
- b) n-Butyllithium, 1.6M (0.0203 mol) was added dropwise at -70°C under N₂ flow to a solution of 1-methyl-1*H*-imidazole (0.0203 mol) in THF (60ml). The mixture was stirred at -70°C for 40 minutes. Chlorotriethylsilane (0.203 mol) was added quickly and the mixture was allowed to warm to 0°C on an ice bath. The mixture was cooled to -70°C and n-butyllithium (0.0203 mol) was added dropwise. The mixture was allowed to

- warm to -20°C and cooled to -70°C. A solution of 2-[3-chloro-4-(4-chlorobenzoyl)-phenyl]-1,2,4-triazine-3,5(2*H*,4*H*)-dione (0.00812 mol) in THF (20ml) was added dropwise. The mixture was allowed to warm to -5°C, then poured out into a saturated NH₄Cl solution and ice, and extracted with EtOAc. The organic layer was separated,
5 dried, filtered and the solvent was evaporated. The residue was purified by column chromatography over silica gel (eluent: CH₂Cl₂/CH₃OH 96/4). The pure fractions were collected and the solvent was evaporated. The residue (0.85g) was crystallized from 2-propanone and diethyl ether. The precipitate was filtered off and dried, yielding 0.47g (13%) of (±)-2-[3-chloro-4-[(chlorophenyl)hydroxy(1-methyl-1*H*-imidazol-5-yl)methyl]phenyl]-1,2,4-triazine-3,5(2*H*,4*H*)-dione monohydrate (compound 121).
10 c) *n*-Butyllithium (0.1 mol) was added dropwise at -70°C under N₂ flow to a solution of *N,N*-dimethylethanamine (0.1 mol) in THF (100ml). The mixture was stirred at -20°C for 30 minutes and cooled again to -70°C. Acetonitrile (0.1 mol) was added dropwise. The mixture was stirred at -20°C for 1 hour and cooled again to -70°C. A solution of
15 2-[3,5-dichloro-4-(4-chlorobenzoyl)phenyl]-1,2,4-triazine-3,5(2*H*,4*H*)-dione (0.05 mol) in THF (100ml) was added dropwise. The mixture was stirred at -70°C for 1 hour, then poured out into NH₄Cl 10% and extracted with EtOAc. The organic layer was separated, dried, filtered and the solvent was evaporated. The residue was purified by column chromatography over silica gel (eluent : CH₂Cl₂/CH₃OH 98/2). Two pure
20 fractions were collected and their solvents were evaporated, yielding 1.62g (8%) of (±)-2,6-dichloro-α-(4-chlorophenyl)-4-[4,5-dihydro-3,5-dioxo-1,2,4-triazin-2(3*H*)-yl]-α-hydroxybenzenepropanenitrile (compound 122).

Example B5

- a) A mixture of intermediate (25) (0.0289 mol) in 2-mercaptoacetic acid (15ml) was
25 stirred at 150°C for 3 hours and then cooled. The mixture was poured out in water, neutralized, and extracted with EtOAc. The organic layer was separated, dried, filtered and the solvent was evaporated. The residue was purified by column chromatography (eluent: CH₂Cl₂/CH₃OH 98/2). The pure fractions were collected and the solvent was evaporated. A sample of this product was crystallized from 2-propanone and diethyl
30 ether. The precipitate was filtered off and dried, yielding 1.2g (±)-2-[3-chloro-4-[(4-chlorophenyl)(1-methyl-1*H*-imidazol-2-yl)methyl]phenyl]-1,2,4-triazine-3,5(2*H*,4*H*)-dione (compound 123)..
b) (±)-2-[3-chloro-4-[(4-chlorophenyl)-3-pyridinylmethyl]phenyl]-1,2,4-triazine-3,5(2*H*,4*H*)-dione was prepared according to the procedure described in example B5a
35 but using 1,2-dimethoxyethane instead of 2-mercaptoacetic acid (compound 124).

Example B6

- a) A mixture of intermediate (50) (0.0027 mol) in toluene (100ml) was stirred and refluxed using a Dean-Stark apparatus. The mixture was decanted and the solvent evaporated. The residue was purified by column chromatography over silica gel (eluent: $\text{CH}_2\text{Cl}_2/\text{CH}_3\text{OH}$ 99/1). The pure fractions were collected and the solvent was evaporated. The residue was crystallized from diethyl ether. The precipitate was filtered off and dried, yielding 1.04g (78%) (\pm)-2-[(3-chloro-4-[(4-chlorophenyl)(3-phenyl-1,2,4-oxadiazol-5-yl)methyl]phenyl]-1,2,4-triazine-3,5(2*H*,4*H*)-dione (compound 125).
- 10 b) (\pm)-2-[3,5-dichloro-4-[(4-chlorophenyl)(3-phenyl-1,2,4-oxadiazol-5-yl)methyl]phenyl]-1,2,4-triazine-3,5(2*H*,4*H*)-dione (comp. 429; mp. 128°C) was prepared analogous to the procedure described in example B6.a except that the starting product was mixed with *p*-toluenesulfonic acid and dimethylsulfoxide instead of toluene.

Example B7

- 15 A mixture of intermediate (66) (0.022 mol) and sodium methoxide, 30% in methanol (0.033 mol) in 1-butanol (350ml) was stirred at RT for 30 minutes. Molecular sieves (12.6g) and then EtOAc (0.033 mol) were added. The mixture was stirred and refluxed overnight, filtered over celite and the solvent was evaporated. The residue was taken up in CH_2Cl_2 , washed with HCl 3N and then with water, dried, filtered and the solvent was
- 20 evaporated. The residue was purified by column chromatography over silica gel (eluent: $\text{CH}_2\text{Cl}_2/\text{CH}_3\text{OH}$ 99/1). The pure fractions were collected and the solvent was evaporated. The residue was crystallized from 2-propanone and DIPE. The precipitate was filtered off and dried, yielding 2.1g (\pm)-2-[3-chloro-4-[(4-chlorophenyl)(5-methyl-1,2,4-oxadiazol-3-yl)methyl]phenyl]-1,2,4-triazine-3,5(2*H*,4*H*)-dione (compound 126).

Example B8

- 25 a) Intermediate (54) (0.00294 mol) was added portionwise at 5°C to phosphoryl chloride (15ml). The mixture was allowed to warm to RT, then stirred at 80°C overnight and cooled. The solvent was evaporated. Ice water was added and the mixture was extracted with CH_2Cl_2 . The organic layer was separated, washed with
- 30 H_2O , dried, filtered and the solvent was evaporated. The residue was purified by column chromatography over silica gel (eluent: $\text{CH}_2\text{Cl}_2/\text{CH}_3\text{OH}$ 97.5/2.5). The pure fractions were collected and the solvent was evaporated. The residue was crystallized from 2-propanone and diethyl ether. The precipitate was filtered off and dried, yielding 0.5g (\pm)-2-[3,5-dichloro-4-[(4-chlorophenyl)(5-phenyl-1,3,4-oxadiazol-2-yl)methyl]-phenyl]-1,2,4-triazine-3,5(2*H*,4*H*)-dione (compound 127).
- 35 b) Compound 127 (0.0114 mol) was dissolved in hexane/ethanol/methanol 50/25/25

(400 ml), then separated into its enantiomers by chiral column chromatography over a Chiralpak AS column (230 g, 20 μ m, I.D.: 5 cm; eluent: hexane/ethanol + 0.1% CF₃COOH/methanol 66/17/17). Two fraction groups were collected. Fraction 1 was added to water. The organic solvent was evaporated and the aqueous concentrate was extracted with CH₂Cl₂. The solvent of the separated organic phase was evaporated. Fraction 2 was treated analogously. Both residues, each individually, were post-purified over Lichroprep 200 (eluent gradient: CH₂Cl₂/CH₃OH). Two pure fraction groups were collected and the solvent was evaporated, yielding 2.86 g (A)-2-[3,5-dichloro-4-[(4-chlorophenyl)(5-phenyl-1,3,4-oxadizol-2-yl)methyl]phenyl]-1,2,4-triazine-3,5(2*H*,4*H*)-dione (compound 189; $\alpha_{20}^D = +50.98^\circ$ ($c = 24.42$ mg/5 ml in CH₃OH)) and 1.75 g (B)-2-[3,5-dichloro-4-[(4-chlorophenyl)(5-phenyl-1,3,4-oxadizol-2-yl)methyl]phenyl]-1,2,4-triazine-3,5(2*H*,4*H*)-dione (compound 190; $\alpha_{20}^D = -50.83^\circ$ ($c = 22.92$ mg/5 ml in CH₃OH)).

Example B9

A mixture of intermediate (59) (0.0108 mol) in toluene (120ml) and methanesulfonic acid (1.05ml) was stirred and refluxed for 4 hours, cooled, poured out into water, decanted, and basified to pH=8 with NH₄OH, while stirring. The aqueous layer was neutralized and extracted with CH₂Cl₂. The organic layer was washed with water, dried, filtered, and the solvent was evaporated. The residue was purified by column chromatography over silica gel (eluent: CH₂Cl₂/CH₃OH 98/2). The desired fractions were collected and the solvent was evaporated. The residue was repurified by HPLC (eluent: CH₃OH/H₂O 80/20). Two pure fractions were collected and their solvents were evaporated, yielding 0.44g (8%) of (\pm)-2-[3-chloro-4-[(4-chlorophenyl)-[5-(phenylamino)-1,3,4-thiadizol-2-yl)methyl]phenyl]-1,2,4-triazine-3,5(2*H*,4*H*)-dione (compound 129), and 0.27g (5%) of (\pm)-2-[3-chloro-4-[(4-chlorophenyl) (4,5-dihydro-4-phenyl-5-thioxo-1*H*-1,2,4-triazol-3-yl)methyl]phenyl]-1,2,4-triazine-3,5(2*H*,4*H*)-dione (compound 128)

Example B10

a) A mixture of intermediate (65) (0.00275 mol) and triethylamine (0.003 mol) in THF (20ml) was stirred at RT. Benzoyl chloride (0.00275 mol) in THF (10ml) was added dropwise and the reaction mixture was stirred at RT for 3 hours. The solvent was evaporated. The residue was stirred in H₂O and CH₂Cl₂. The organic layer was dried, filtered, and the solvent was evaporated. The residue was purified over silica gel on a glass filter (eluent: CH₂Cl₂/CH₃OH 98/2). The desired fractions were collected and the solvent was evaporated. The residue was repurified by column chromatography over silica gel (eluent: CH₂Cl₂/CH₃OH 98/2). The desired fractions were collected and the

solvent was evaporated. The residue was stirred in DIPE, filtered off, washed with DIPE and dried. The residue was repurified by HPLC over silica gel (eluent: $\text{CH}_2\text{Cl}_2/\text{CH}_3\text{OH}$ 98/2). The desired fractions were collected and the solvent was evaporated. The residue was stirred in DIPE. The precipitate was filtered off, washed and dried, yielding 0.4 g (\pm)-*N*-[[2-chloro-4-(4,5-dihydro-3,5-dioxo-1,2,4-triazin-3(2*H*)-yl)phenyl](4-chlorophenyl)methyl]benzamide (compound 47).

5 b) A mixture of intermediate 65 (0.00275 mol) and 2-methylthiothiazolo[5,4-b]pyridine (0.0035 mol) was heated up to 170 °C and stirred for 2 days. The reaction mixture was dissolved in $\text{CH}_2\text{Cl}_2/\text{CH}_3\text{OH}$ (90/10). The precipitate was filtered off and the filtrate was evaporated. The residue was purified by flash column chromatography over silica gel (eluent: $\text{CH}_2\text{Cl}_2/\text{CH}_3\text{OH}$ 99/1). The desired fractions were collected and the solvent was evaporated. The residue was stirred in DIPE, filtered off, washed and dried, yielding 0.1 g of (\pm)-2-[3-chloro-4-[(4-chlorophenyl)[(thiazolo[5,4-b]pyridin-2-yl)-amino]methyl]phenyl]-1,2,4-triazine-3,5(2*H*,4*H*)-dione. (comp. 410)

15 Example B11

A solution of intermediate (63) (0.0080 mol), 6-chloro-2,4-dimethoxypyrimidine (0.0084 mol) and 1,8-diazabicyclo[5.4.0]undec-7-ene (0.0088 mol) in DMF (50 ml) was stirred for 4 days at RT. The solvent was evaporated and the residue was stirred in water and this mixture was extracted with $\text{CH}_2\text{Cl}_2/\text{CH}_3\text{OH}$ 90/10. The separated organic layer was dried, filtered, and the solvent was evaporated. The residue was purified by column chromatography over silica gel (eluent: $\text{CH}_2\text{Cl}_2/\text{CH}_3\text{OH}$ 98/2). The desired fractions were collected and the solvent was evaporated. The residue was repurified by reversed-phase liquid chromatography over silica gel (eluent: (0.5% NH_4OAc in H_2O)/ $\text{CH}_3\text{OH}/\text{CH}_3\text{CN}$ 28/36/36, upgrading to 0/50/50). The desired fractions were collected and the solvent was evaporated. The residue was stirred in DIPE, filtered off, washed, then dried, yielding 0.4 g (\pm)-2-[3-chloro-4-[(4-chlorophenyl)[(2,6-dimethoxy-2-pyrimidinyl)thio]methyl]phenyl]-1,2,4-triazine-3,5(2*H*,4*H*)-dione (compound 96).

Example B12

30 A solution of intermediate (48) (0.012 mol) in pyridine (45ml) was added to a solution of 2-mercapto-2-benzenamine (0.0132 mol) in pyridine (30ml). The mixture was stirred and heated at 60°C for 18 hours, poured out into HCl 3N, and extracted with CH_2Cl_2 . The organic layer was separated, washed with H_2O , dried, filtered and the solvent was evaporated. The residue was purified by column chromatography over silica gel (eluent: $\text{CH}_2\text{Cl}_2/\text{CH}_3\text{OH}$ 99/1). The pure fractions were collected and the solvent was

evaporated, yielding 1.23 g (21%) (\pm)-2-[4-[2-benzothiazolyl-(4-chlorophenyl)methyl]-3-chlorophenyl]-1,2,4-triazine-3,5(2*H*,4*H*)-dione (compound 130).

Example B13

5 a) A mixture of 2,6-dichloro- α -(4-chlorophenyl)-4-(4,5-dihydro-3,5-dioxo-1,2,4-triazin-2(3*H*)-yl)benzeneethanethionate (0.00453 mol) and 2-bromo-1-phenylethanone (0.00498 mol) in ethanol (80ml) was stirred and refluxed for 2 hours. The solvent was evaporated. The residue was taken up in CH_2Cl_2 , washed with K_2CO_3 10% and then with water, dried, filtered and the solvent was evaporated. The residue was purified by column chromatography over silica gel (eluent: $\text{CH}_2\text{Cl}_2/\text{CH}_3\text{OH}/\text{NH}_4\text{OH}$ 90/10/0.5;).

10 The pure fractions were collected and the solvent was evaporated. The residue was crystallized from DIPE. The precipitate was filtered off and dried, yielding 1.05 g (43%) of (\pm)-2-[3,5-dichloro-4-[(4-chlorophenyl)(4-phenyl-2-thiazolyl)methyl]phenyl]-1,2,4-triazine-3,5(2*H*,4*H*)-dione (compound 38).

15 b) (\pm)-2-[3,5-dichloro-4-[(4-chlorophenyl)[5-(1-methylethyl)-4-phenyl-2-thiazolyl)methyl]phenyl]-1,2,4-triazine-3,5(2*H*,4*H*)-dione (comp. 241) was prepared according to example B13.a and in addition triethylamine was used as a base.

c) 2,6-Dichloro- α -(4-chlorophenyl)-4-(4,5-dihydro-3,5-dioxo-1,2,4-triazin-2(3*H*)-yl)benzeneethanethioamide (0.031 mol) was added at RT to a solution of (\pm)-1,1-dimethylethyl α -bromo- β -oxo-benzenepropanoate (0.0465 mol) and K_2CO_3 (0.093 mol) in 20 CH_3CN (190ml). The mixture was stirred at RT for 3.5 hours. H_2O was added. The mixture was acidified with HCl 3*N* and extracted with EtOAc . The organic layer was separated, dried, filtered and the solvent was evaporated. The residue was purified by column chromatography over silica gel (eluent: $\text{CH}_2\text{Cl}_2/\text{CH}_3\text{OH}$ 99/1). The pure fractions were collected and the solvent was evaporated, yielding 11 g (54%) of (\pm)-1,1-25 dimethylethyl 2-[(4-chlorophenyl)[2,6-dichloro-4-(4,5-dihydro-3,5-dioxo-1,2,4-triazin-2(3*H*)-yl)phenyl)methyl]-4-phenyl-5-thiazolcarboxylate (comp. 298).

Example B14

A mixture of 2,6-dichloro- α -(4-chlorophenyl)-4-(4,5-dihydro-3,5-dioxo-1,2,4-triazin-2(3*H*)-yl)benzeneethanethionate (0.0197 mol) and 1-bromo-2,2-diethoxyethane 30 (0.0256 mol) in HCl 3*N* (10ml) and ethanol (145ml) was stirred and refluxed for 5 hours. The solvent was evaporated. The residue was taken up in CH_2Cl_2 , washed with K_2CO_3 10% and extracted with CH_2Cl_2 . The organic layer was separated, washed with water, dried, filtered and the solvent was evaporated. The residue was purified by column chromatography over silica gel (eluent: $\text{CH}_2\text{Cl}_2/\text{CH}_3\text{OH}/\text{NH}_4\text{OH}$ 85/15/1). The 35 pure fractions were collected and the solvent was evaporated. The residue was crystallized from DIPE. The precipitate was filtered off, dried and recrystallized from

2-propanone and diethyl ether. The precipitate was filtered off and dried, yielding 1.32g (\pm)-2-[3,5-dichloro-4-[(4-chlorophenyl)-2-thiazolylmethyl]-phenyl]-1,2,4-triazine-3,5(2*H*,4*H*)-dione (compound 39).

Example B15

- 5 a) A mixture of intermediate (52) (0.0076 mol) in EtOAc (45ml) was stirred and refluxed for 18 hours, then poured out into H₂O and extracted with EtOAc. The organic layer was separated, washed with K₂CO₃ 10% and with H₂O, dried, filtered and the solvent was evaporated. The residue was purified by column chromatography over silica gel (eluent: CH₂Cl₂/CH₃OH 99/1). The pure fractions were collected and the
- 10 solvent was evaporated, yielding 0.9g (25%) of (\pm)-2-[4-[2-benzoxazolyl(4-chlorophenyl)methyl]-3-chlorophenyl]-1,2,4-triazine-3,5(2*H*,4*H*)-dione (compound 131).
- b) (\pm)-2-[3-chloro-4-[1-(4-chlorophenyl)-1-(2-benzoxazolyl)ethyl]phenyl]-1,2,4-triazine-3,5(2*H*,4*H*)-dione was prepared using the same procedure as in example B15a but by using methanesulfonic acid instead of acetic acid (compound 132).

Example B16

- 15 A mixture of compound (33) (0.0231 mol) in methanol (100ml) and sulfonic acid (2ml) was stirred and refluxed for 3 days, then cooled, poured out on ice, neutralized and extracted with CH₂Cl₂. The organic layer was separated, dried, filtered and the solvent was evaporated. The residue was purified by column chromatography over silica gel
- 20 (eluent: CH₂Cl₂/CH₃OH 99/1 to 98/2). The pure fractions were collected and the solvent was evaporated, yielding 4 g (38%) of (\pm)-2-[3-chloro-4-[(4-chlorophenyl)-methoxy(2-thiazolyl)methyl]phenyl]-1,2,4-triazine-3,5(2*H*,4*H*)-dione (compound 37).

Example B17

- a) Compound (33) (0.00425 mol) was dissolved in thionyl chloride (20ml) at 10°C, and
- 25 the mixture was stirred at RT for 4 hours. The solvent was evaporated, yielding (\pm)-2-[3-chloro-4-[chloro(4-chlorophenyl)-2-thiazolylmethyl]phenyl]-1,2,4-triazine-3,5(2*H*,4*H*)-dione (compound 36).
- b) A solution of compound (36) (0.00425 mol) in THF (20ml) was added dropwise at 5°C to NH₄OH (20ml) and the mixture was stirred at RT for 2 hours, then poured out
- 30 on ice, neutralized with HCl 6N and extracted with EtOAc. The organic layer was separated, dried, filtered and the solvent was evaporated. The residue was purified by column chromatography over silica gel (eluent: CH₂Cl₂/CH₃OH 97/3). The pure fractions were collected and the solvent was evaporated. The residue was repurified by column chromatography over Kromasil C18 (eluent: CH₃OH/H₂O/HOAc 70/30/1). The
- 35 pure fractions were collected and the solvent was evaporated. The residue was taken up in H₂O and NH₄OH (pH=8) was added. The precipitate was filtered off, washed with

H₂O and diethyl ether, and dried, yielding 0.3g (±)-2-[3-chloro-4-[amino-(4-chloro-phenyl)-2-thiazolylmethyl] phenyl]-1,2,4-triazine-3,5(2*H*,4*H*)-dione (compound 35).

Example B.18

A mixture of 2-[3,5-dichloro-4-[(4-chlorophenyl)hydroxymethyl]phenyl]-1,2,4-triazine-3,5(2*H*,4*H*)-dione (0.005 mol) and 5-phenyl-1,3,4-oxadiazole-2(3*H*)-thione (0.006 mol) in methanesulfonic acid (20 ml) was stirred for 18 hours at RT. The reaction mixture was poured out into water/ice (150 ml), and the resulting precipitate was filtered off, stirred in water, treated with NaHCO₃ and this mixture was extracted with CH₂Cl₂. The separated organic layer was dried, filtered and the solvent evaporated. The residue was purified by flash column chromatography over silica gel (eluent: CH₂Cl₂/CH₃OH 97.5/2.5). The pure fractions were collected and the solvent was evaporated. The residue was stirred in DIPE, filtered off and dried, yielding 1 g of (±)-2-[3,5-dichloro-4-[(4-chlorophenyl)[(5-phenyl-1,3,4-oxadiazol-2-yl)thio]methyl]phenyl]-1,2,4-triazine-3,5(2*H*,4*H*)-dione (comp. 406).

Example B.19

- a) A solution of 2,6-dichloro- α -(4-chlorophenyl)-4-(4,5-dihydro-3,5-dioxo-1,2,4-triazin-2(3*H*)-yl)benzeneacetyl chloride (0.188 mol) in 1,4-dioxane (900 ml) was stirred at RT. NaBH₄ (36.25 g) was added portionwise over 2.5 hours. The resulting reaction mixture was stirred for 3 hours at RT. The reaction mixture was cooled and acidified till pH 6 with 1 N HCl. The precipitated salts were removed by filtration. The filtrate was washed with water, and the precipitate was filtered off, stirred in DIPE, filtered off and dried, yielding 22.5 g of (±)-2-[3,5-dichloro-4-[1-(4-chlorophenyl)-2-hydroxyethyl]phenyl]-1,2,4-triazine-3,5(2*H*,4*H*)-dione (comp. 420). The biphasic filtrate was separated into its layers. The organic layer was dried, filtered and the solvent was evaporated. The residue was purified by HPLC over silica gel (eluent: CH₂Cl₂/CH₃OH 98.5/1.5 and 97/3). The pure fractions were collected and the solvent was evaporated. The residue was stirred in DIPE, filtered off, washed, and dried, yielding 24 g of (±)-2-[3,5-dichloro-4-[1-(4-chlorophenyl)-2-hydroxyethyl]phenyl]-1,2,4-triazine-3,5(2*H*,4*H*)-dione (comp. 420).
- b) A solution of compound 420 (0.01 mol) and *N*-ethyl-*N*-(1-methylethyl)-2-propanamine (0.02 mol) in 1,4-dioxane (80 ml) was stirred at 5-10°C under N₂ atmosphere. A solution of methanesulfonyl chloride (0.02 mol) in 1,4-dioxane (10 ml) was added dropwise at 5-10 °C. The resulting reaction mixture was stirred for one hour at RT. The solvent was evaporated under reduced pressure. The residue was dissolved in CH₂Cl₂, washed with water, dried, filtered and the solvent was evaporated, yielding 4.9 g of (±)-2-[2,6-dichloro-4-(4,5-dihydro-3,5-dioxo-1,2,4-triazin-2(3*H*)-yl)phenyl]-2-(4-chlorophenyl)ethanol methanesulfonate (ester) (comp. 435).

c) A mixture of compound 435 (0.001 mol), 2-pyridinethiol (0.0012 mol) and NaHCO₃ (0.0012 mol) in DMF (30ml) was stirred at RT under N₂ flow, then heated to 60°C and stirred for 48 hours. 2-pyridinethiol (0.0012 mol) and NaHCO₃ (0.0012 mol) were added again. The mixture was stirred for 1 day. 2-pyridinethiol (0.006 mol) was added again and the mixture was stirred and refluxed for 1 day. The solvent was evaporated under reduced pressure. The residue was dissolved in CH₂Cl₂ and extracted with water. The organic layer was separated, dried, filtered and the solvent was evaporated. The residue was purified by HPLC (eluent: NH₄OAc 0.5% in H₂O/CH₃OH/CH₃CN 67.5/7.5/25 to 0/50/50 after 10 minutes to 0/0/100 after 10 minutes). The desired fractions were collected and the solvent was evaporated. The residue was stirred in DIPE. The precipitate was filtered off, washed and dried, yielding 0.05g (10%) of (±)-2-[3,5-dichloro-4-[1-(4-chlorophenyl)-2-(2-pyridinylthio)ethyl]phenyl]-1,2,4-triazine-3,5(2*H*,4*H*)-dione (comp. 422).

Example B.20

A mixture of intermediate 75 (0.0159 mol) in dimethylsulfoxide (170ml) and H₂O (20ml) was stirred at 160°C for 3 hours. The mixture was cooled and poured out on ice. The precipitate was filtered off, washed with H₂O and taken up in EtOAc. The organic solution was dried, filtered and the solvent was evaporated. The residue was purified by column chromatography over silica gel (eluent: CH₂Cl₂/CH₃OH 99.5/0.5 to 96/4). The desired fraction was collected and the solvent was evaporated, yielding 2015g of (±)-2-[3-chloro-4-[(4-chlorophenyl)(4-phenyl-2-thiazolyl)methyl]phenyl]-1,2,4-triazine-3,5(2*H*,4*H*)-dione (comp. 419; mp. 90°C).

Example B.21

A mixture of (±)-2-[4-[3-bromo-1-(4-chlorophenyl)-2-oxopropyl]-3,5-dichlorophenyl]-1,2,4-triazine-3,5(2*H*,4*H*)-dione (0.0025 mol) and benzenecarbothioamide (0.0025 mol) in ethanol (25 ml) was stirred and refluxed for 3 hours, then stirred overnight at RT. The solvent was evaporated. The residue was purified twice by column chromatography over silica gel (eluent: CH₂Cl₂/CH₃OH (1) 97/3 and (2) 98/2 v/v). The desired fractions were collected and the solvent was evaporated. The residue was repurified by column chromatography over silica gel (eluent: CH₂Cl₂/CH₃OH 99/1). The pure fractions were collected and the solvent was evaporated. The residue was stirred in hexane, filtered off, then dried, yielding 0.3 g (22%) of (±)-2-[3,5-dichloro-4-[(4-chlorophenyl)(2-phenyl-4-thiazolyl)methyl]phenyl]-1,2,4-triazine-3,5(2*H*,4*H*)-dione (comp. 363).

Example B.22

a) Compound 298 (0.0137 mol) was added at 10°C under N₂ flow to trifluoroacetic acid (120ml). The mixture was allowed to warm to RT and stirred for 1 hour. H₂O was

- added. The precipitate was filtered off, washed with H₂O and taken up in CH₂Cl₂ and a small amount of CH₃OH. The organic layer was separated, dried, filtered and the solvent was evaporated. The residue was crystallized from CH₃CN. The precipitate was filtered off and the filtrate was evaporated. The residue was purified by column chromatography over silica gel (eluent: CH₂Cl₂/CH₃OH/HOAc 97/3/0.1). The pure fractions were collected and the solvent was evaporated. This fraction was crystallized from CH₃CN. The precipitate was filtered off and dried, yielding 1.34g (67%) of (±)-2-[(4-chlorophenyl)[2,6-dichloro-4-(4,5-dihydro-3,5-dioxo-1,2,4-triazin-2(3H)-yl)phenyl]methyl]-4-phenyl-5-thiazolcarboxylic acid (Comp. 299; mp 206°C).
- 10 b) 1,1'-Carbonylbis-1H-imidazole (0.0081 mol) was added to a suspension of compound 299 (0.00324 mol) in CH₂Cl₂ (25ml). The mixture was stirred at RT for 2 hours. Dimethylamine (0.00324 mol) was added. The mixture was stirred at RT for 48 hours. H₂O was added. The mixture was acidified with HCl 3N and extracted with CH₂Cl₂. The organic layer was separated, dried, filtered and the solvent was evaporated.
- 15 The residue was purified by column chromatography over silica gel (eluent: CH₂Cl₂/CH₃OH 98.5/1.5). The pure fractions were collected and the solvent was evaporated. The residue was crystallized from DIPE. The precipitate was filtered off and dried, yielding 1.04g (52%) of (±)-N,N-dimethyl-2-[(4-chlorophenyl)[2,6-dichloro-4-(4,5-dihydro-3,5-dioxo-1,2,4-triazin-2(3H)-yl)phenyl]methyl]-4-phenylthiazol-5-carboxamide
- 20 (Comp. 303; mp 150°C).

Example B.23

- a) A solution of compound 350 (0.014 mol) in 2,6-dimethylpyridine (1.63 ml) and THF (80 ml) was stirred and cooled to -78 °C. Trifluoromethanesulfonic anhydride (0.014 mol) was added dropwise and the mixture was stirred for 7 hours at -78 °C,
- 25 yielding (±)-2-[(4-chlorophenyl)[2,6-dichloro-4-(4,5-dihydro-3,5-dioxo-1,2,4-triazin-2(3H)-yl)phenyl]methyl]thio]-4-pyrimidinol trifluoromethanesulfonate (ester) (comp. 356).
- b) A mixture of compound 356 (0.0047 mol) in THF (35 ml) was stirred at RT. 2-Aminoethanol (0.0235 mol) was added. The reaction mixture was stirred for one hour at 50 °C, then for 16 hours at RT. The solvent was evaporated. The residue was
- 30 purified by flash column chromatography over silica gel (eluent: CH₂Cl₂/CH₃OH 99/1, 98/2 and 93/7). The desired fractions were collected and the solvent was evaporated. The residue was repurified by HPLC over silica gel (eluent: CH₂Cl₂/CH₃OH from 100/0 over 30 minutes to 92/8). The desired fractions were collected and the solvent was evaporated. The residue was stirred in DIPE, filtered off, washed and dried, yielding
- 35 0.3 g of (±)-2-[3,5-dichloro-4-[(4-chlorophenyl)[4-[(2-hydroxyethyl)amino]-2-pyrimidinyl]-thio]methyl]phenyl]-1,2,4-triazine-3,5(2H,4H)-dione (comp. 357).

Example B.24

- a) LiCl (0.035 mol) was added portionwise at 80°C to a mixture of compound 285 (0.007 mol) and KBH₄ (0.035 mol) in THF (45ml). The mixture was stirred at 80°C for 4 hours. KBH₄ (0.035 mol) and then LiCl (0.035 mol) were added. The mixture was stirred at 80°C for 4 hours, at RT overnight, then poured out into ice water, acidified with HCl 3N and extracted with CH₂Cl₂. The organic layer was separated, dried, filtered and the solvent was evaporated. The residue was purified by column chromatography over silica gel (eluent: CH₂Cl₂/CH₃OH 97/3; 20-45 µm). The pure fractions were collected and the solvent was evaporated, yielding 2.1g (51%) of (±)-2-[3,5-dichloro-4-
5 [(4-chlorophenyl)[4-(2-fluorophenyl)-5-(hydroxymethyl)-2-thiazolyl]methyl]phenyl]-1,2,4-triazine-3,5(2*H*,4*H*)-dione (comp. 323).
- b) Thionylchloride (0.0113 mol) was added at 10°C to a mixture of compound 323 (0.0094 mol) in CH₂Cl₂ (30 ml). The mixture was stirred at RT for 2.5 hours, washed with H₂O and with K₂CO₃ 10%, dried, filtered and the solvent was evaporated, yielding
15 2g of (±)-2-[4-[[5-(chloromethyl)-4-(2-fluorophenyl)-2-thiazolyl](4-chlorophenyl)methyl]-3,5-dichlorophenyl]-1,2,4-triazine-3,5(2*H*,4*H*)-dione (comp. 324).
- c) A mixture of compound 324 (0.0034 mol), dimethylamine (0.0068 mol) and K₂CO₃ (0.0102 mol) in CH₃CN (100ml) was stirred and refluxed for 3 hours and then cooled. The solvent was evaporated. The residue was taken up in CH₂Cl₂. The organic solution
20 was washed with H₂O, dried, filtered and the solvent was evaporated. The residue was purified by column chromatography over silica gel (eluent: CH₂Cl₂/CH₃OH/H₂O 97/3/0.4). The pure fractions were collected and the solvent was evaporated. The residue was crystallized from diethyl ether and CH₃CN. The precipitate was filtered off and dried, yielding 0.84g of (±)-2-[4-[(4-chlorophenyl)[5-[(dimethylamino)methyl]-4-(2-
25 fluorophenyl)-2-thiazolyl]methyl]-3,5-dichlorophenyl]-1,2,4-triazine-3,5(2*H*,4*H*)-dione (comp. 325; mp 250°C).

Example B.25

- A mixture of compound 229 (0.0041 mol) and triethylamine (0.0082 mol) in CH₂Cl₂ (45ml) was stirred at RT for 1 hour. A solution of acetyl chloride (0.0041 mol) in
30 CH₂Cl₂ (5ml) was added at 10°C. The mixture was stirred at RT for 12 hours, then poured out into H₂O and decanted. The organic layer was washed with HCl 3N and with H₂O, dried, filtered and the solvent was evaporated. The residue was purified by column chromatography over silica gel (eluent: CH₂Cl₂/CH₃OH 95/5). The pure fractions were collected and the solvent was evaporated. The residue was crystallized
35 from CH₃CN and DIPE. The precipitate was filtered off and dried, yielding 0.52g of (±)-2-[3,5-dichloro-4-[(4-chlorophenyl)[4-(4-piperidiny)-2-thiazolyl]methyl]phenyl]-1,2,4-triazine-3,5-(2*H*,4*H*)-dione monohydrochloride (comp. 230; mp 212°C).

Example B.26

A mixture of compound 212 (0.00646 mol) in $\text{NH}_3/\text{CH}_3\text{OH}$ 7N (100ml) was stirred and refluxed for 3 hours and then cooled. The solvent was evaporated. The residue was taken up in EtOAc and a small amount of CH_3OH . The organic layer was separated,
5 washed with HCl 3N, dried, filtered and the solvent was evaporated. The residue was purified by column chromatography over silica gel (eluent: $\text{CH}_2\text{Cl}_2/\text{CH}_3\text{OH}$ 97/3). The pure fractions were collected and the solvent was evaporated. The residue was crystallized from 2-propanone and diethyl ether. The precipitate was filtered off and dried, yielding 0.85g of (\pm)-*N*-[2-[5-[(4-chlorophenyl)[2,6-dichloro-4-(4,5-dihydro-3,5-
10 dioxo-1,2,4-triazin-2(3*H*)-yl)phenyl]methyl]-1,3,4-oxadiazol-2-yl]phenyl]-2-hydroxyacetamide (comp. 213; mp 235°C).

Example B.27

A mixture of compound 352 (0.005 mol) in HBr (75 ml; 48%) was stirred at RT. The mixture was warmed to 140°C on an oil bath and stirred for 30 minutes. The mixture
15 was cooled. The solvent was evaporated. H_2O was added. The mixture was neutralized with NaOH 50% and extracted with CH_2Cl_2 . The product was filtered off and stirred in CH_3OH , in CH_3CN and then in CH_2Cl_2 , and dried. This fraction was stirred in H_2O (20 ml), and CH_3COOH (± 1 equiv) was added. The product was filtered off, washed with H_2O and dried, yielding 1.3 g of (\pm)-2-[3,5-dichloro-4-[(4-chlorophenyl)[4-(1-piperazinyl)-2-
20 pyrimidinyl]thio]methyl]phenyl]-1,2,4-triazine-3,5(2*H*,4*H*)-dione monohydrate (comp. 360).

Example B.28

a) A mixture of compound 192 (0.014 mol) in THF (100 ml) and methanol (100 ml) was hydrogenated at 50°C with platina on activated charcoal (2 g; 10%) as a catalyst in the presence of a thiophene solution (2 ml). After uptake of H_2 , the catalyst was filtered
25 off and the filtrate was evaporated. Toluene was added and azeotroped on the rotary evaporator, yielding 6.2 g of (\pm)-2-[4-[[5-(3-aminophenyl)-1,3,4-oxadiazol-2-yl](4-chlorophenyl)methyl]-3,5-dichlorophenyl]-1,2,4-triazine-3,5(2*H*,4*H*)-dione (comp. 193).
b) Compound 193 (0.012 mol) was dissolved in acetic acid (40 ml) and HCl (3.6 ml) at about 5°C. A solution of NaNO_2 (0.0126 mol) in H_2O (10 ml) was added dropwise at
30 5°C. The reaction mixture was stirred for 1 hour at 5 °C. NaN_3 (0.0126 mol) was added portionwise. The reaction mixture was stirred for 30 minutes, then poured out onto ice. The precipitate was filtered off, washed with water, then dissolved in CH_2Cl_2 . The organic solution was dried, filtered, and the solvent was evaporated. The residue was purified by HPLC over silica gel (eluent: $\text{CH}_2\text{Cl}_2/\text{CH}_3\text{OH}$ 98/2). The pure fractions were
35 collected and the solvent was evaporated. The residue was stirred in boiling ethanol, filtered off and washed with ethanol/DIPE, then dried, yielding 2.1 g of (\pm)-2-[4-[[5-(3-azidophenyl)-1,3,4-oxadiazol-2-yl](4-chlorophenyl)methyl]-3,5-dichlorophenyl]-1,2,4-

triazine-3,5(2*H*,4*H*)-dione (comp. 194).

Example B.29

- a) A mixture of compound 328 (0.00271 mol) in HBr (20ml; 33% in HOAc) and HBr (20ml; 48% in H₂O) was stirred and refluxed overnight, then cooled, poured out into ice water, neutralized with a concentrated NaOH solution and centrifuged. The residue was washed with H₂O and dried. The residue was purified by column chromatography over silica gel (eluent: CH₂Cl₂/CH₃OH 95/5). The pure fractions were collected and the solvent was evaporated. The residue was taken up in CH₃OH and CH₂Cl₂. The organic solution was washed with a solution at pH 4 and a solution at pH 7, then dried.
- 10 Activated charcoal was added. The mixture was filtered over celite. The solvent was evaporated. The residue was crystallized from CH₃CN and diethyl ether. The precipitate was filtered off and dried, yielding 0.27g of (±)-2-[4-[[5-(aminomethyl)-4-phenyl-2-thiazolyl](4-chlorophenyl)methyl]-3,5-dichlorophenyl]-1,2,4-triazine-3,5(2*H*,4*H*)-dione (comp. 329; mp 170°C).
- 15 b) A solution of compound 329 (0.0035 mol) and isothiocyanatobenzene (0.0042 mol) in THF (25ml) was stirred at RT for 90 minutes. The solvent was evaporated. The residue was dissolved in CH₂Cl₂. The organic solution was washed with H₂O, dried, filtered and the solvent was evaporated. The residue was purified by column chromatography over silica gel (eluent: CH₂Cl₂/CH₃OH 98/2). The pure fractions were collected
- 20 and the solvent was evaporated. The residue was taken up in DIPE. The precipitate was filtered off and dried, yielding 0.64g (±)-*N*-[[2-[[4-(4-chlorophenyl)[2,6-dichloro-(4,5-dihydro-3,5-dioxo-1,2,4-triazin-2(3*H*)-yl)phenyl]methyl]-4-phenyl-5-thiazolyl]methyl]-*N'*-phenylthiourea (comp. 331; mp 159°C).

Example B.30


- 25 a) TiCl₃ (0.034 mol; 15% aqueous solution) was added dropwise at RT to a mixture of compound 216 (0.0034 mol) in THF (60ml). The mixture was stirred at RT for 5 hours, poured out into H₂O and extracted with EtOAc. The organic layer was separated, washed with H₂O, dried, filtered and the solvent was evaporated, yielding 1.9g of (±)-2-[4-[[5-(3-amino-2-methylphenyl)-1,3,4-oxadiazol-2-yl](4-chlorophenyl)methyl]-3,5-dichlorophenyl]-1,2,4-triazine-3,5(2*H*,4*H*)-dione (comp. 217).
- 30 b) A mixture of (acetyloxy)acetyl chloride (0.0121 mol) in CH₂Cl₂ (15ml) was added at 10°C under N₂ flow to a mixture of compound 217 (0.011 mol) and *N*-ethyl-*N*-(1-methylethyl)-2-propanamine (0.0165 mol) in CH₂Cl₂ (60 ml). The mixture was stirred at RT for 12 hours, poured out into H₂O, acidified with HCl 3N and extracted with
- 35 CH₂Cl₂. The organic layer was separated, dried, filtered and the solvent was evaporated. The residue was purified by column chromatography over silica gel (eluent: CH₂Cl₂/CH₃OH 97/3). The pure fractions were collected and the solvent was

evaporated. Part of the residue (0.9g) was crystallized from diethyl ether and CH₃CN. The precipitate was filtered off and dried, yielding 0.65g of (±)-2-(acetyloxy)-N-[3-[5-[(4-chlorophenyl)[2,6-dichloro-(4,5-dihydro-3,5-dioxo-1,2,4-triazin-2(3H)yl)phenyl]methyl]-1,3,4-oxadiazol-2-yl]-2-methylphenyl]acetamide. (comp. 223; mp 206°C).

5


Tables 1 to 8 list compounds of the present invention as prepared according to one of the above examples. These all are racemic mixtures unless otherwise mentioned.

Table 1




Co. No.	Ex. No.	R ^{5a}	R ^{5b}	R ^{11a}	R ^{11b}	Melting-point °C
134	B5a	Cl	H	phenyl	CH ₃	180°C
135	B5a	Cl	Cl	2-Cl-phenyl	CH ₃	170°C
136	B5a	Cl	Cl	phenyl	CH ₃	175°C
137	B5a	Cl	Cl	CH ₃	2-Cl-phenyl	120°C
138	B5a	Cl	Cl	CH ₃	phenyl	120°C
139	B5a	Cl	H	2-Cl-phenyl	CH ₃	>260°C
140	B5a	Cl	H	phenyl	phenyl	186-188°C
141	B5a	Cl	H	H	phenyl	168°C
142	B5a	Cl	Cl	H	phenyl	
143	B5a	Cl	H	3-F-phenyl	CH ₃	146°C
144	B5a	H	Cl	2-Cl-phenyl	phenyl	140°C
145	B5a	Cl	Cl	2-Cl-phenyl	phenyl	160°C
146	B5a	H	H	phenyl	CH ₃	230°C
147	B5a	Cl	Cl	2-Cl-phenyl	2-Cl-phenyl	158°C
148	B5a	Cl	H	3-F-phenyl	H	155°C
149	B5a	Cl	H	4-Cl-phenyl	CH ₃	145°C
150	B5a	Cl	H	phenyl	2-Cl-phenyl	220°C
151	B5a	Cl	H	2-Cl-phenyl	2-Cl-phenyl	150°C
152	B5a	Cl	Cl	2-F-phenyl	CH ₃	185°C
153	B5a	H	OCH ₃	phenyl	CH ₃	163°C
154	B5a	Cl	H	CH ₃	2-Cl-phenyl	190°C

Table 2




Co. No.	Ex. No.	R ¹	R ^{4a}	R ^{5a}	R ^{11a}	salt form / stereochemistry / melting point
155	B8a	CH ₃	CF ₃	H	phenyl	126°C
156	B8a	H	H	Cl	2-F-phenyl	169°C
157	B8a	H	H	Cl	3-Cl-phenyl	188°C
158	B8a	H	H	Cl	4-pyridinyl	H ₂ O (1:1)/170°C
159	B8a	H	H	Cl	cyclohexyl	164°C
160	B8a	H	H	Cl	3-F-phenyl	156°C
161	B8a	H	H	Cl	2-furanyl	170°C
162	B8a	H	H	Cl	methyl	120°C
163	B8a	H	H	Cl	2-Cl-phenyl	H ₂ O (1:1)/160°C
164	B8a	H	H	Cl	propyl	135°C
165	B8a	H	CF ₃	Cl	phenyl	212°C
166	B8a	H	H	Cl	2-thienyl	180°C
167	B8a	H	H	Cl	4-Cl-phenyl	230°C
168	B8a	H	H	Cl	4-Br-phenyl	
169	B8a	H	H	Cl	2-pyridinyl	182°C
170	B8a	H	H	Cl	3-methoxyphenyl	208°C
171	B8a	H	H	Cl	4-methoxyphenyl	212°C
172	B8a	H	H	Cl	phenylethyl	148°C
173	B8a	H	H	Cl	phenyl-CH ₂ -	190°C
174	B8a	H	H	Cl	2-(methoxy)phenyl	164°C
175	B8a	H	H	Cl	(2-Cl-phenyl)-O-CH ₂ -	135°C
176	B8a	H	H	Cl	C ₂ H ₅ -O-CO-CH ₂ -	177°C
177	B8a	H	H	Cl	4-CH ₃ -phenyl	>260°C
178	B8a	H	H	Cl	3-CH ₃ -phenyl	188°C
179	B8a	H	H	Cl	NC-CH ₂ -	222°C
180	B8a	H	H	Cl	4-[N(CH ₃) ₂]-phenyl	224°C
181	B8a	H	H	Cl	C ₂ H ₅ -O-(CH ₂) ₂ -	130°C
182	B8a	H	H	Cl	3-[N(CH ₃) ₂]-phenyl	240°C
183	B2a	H	H	Cl	4-nitrophenyl	
184	B28	H	H	Cl	4-aminophenyl	
185	B28b	H	H	Cl	4-(-N=N ⁺ =N ⁻)-phenyl	
186	B8a	H	H	Cl	C ₂ H ₅ -O-CO-	137°C
187	B8a	H	H	Cl	phenyl-O-(CH ₂) ₂ -	215°C



Co. No.	Ex. No.	R ¹	R ^{4a}	R ^{5a}	R ^{11a}	salt form / stereochemistry / melting point
188	B8a	H	H	Cl	2-CH ₃ -phenyl	150°C
189	B8b	H	H	Cl	phenyl	(A)
190	B8b	H	H	Cl	phenyl	(B)
191	B8a	H	H	Cl	1-(C ₂ H ₅ -O-CO)-4-piperidinyl	230°C
192	B8a	H	H	Cl	3-nitrophenyl	
193	B28a	H	H	Cl	3-aminophenyl	
194	B28b	H	H	Cl	3-(-N=N ⁺ =N ⁻)-phenyl	
195	B8a	H	H	Cl	1-CH ₃ -4-piperidinyl	
196	B8a	H	H	Cl	1-CH ₃ -3-piperidinyl	150°C
197	B8a	H	H	Cl	Cl-CH ₂ -	
198	B24c	H	H	Cl	(CH ₃) ₂ -N-CH ₂ -	188°C
199	B8a	H	H	Cl	4-(4-CH ₃ -1-piperazinyl)phenyl	150°C
200	B8a	H	H	Cl	3-OH-phenyl	159°C
201	B8a	H	H	Cl	3-pyridinyl	190°C
202	B8a	H	H	Cl	2-hydroxyphenyl	180°C
203	B8a	H	H	Cl	3-CH ₃ -2-thienyl	161°C
204	B8a	H	H	Cl	3-(NH ₂ -SO ₂)-phenyl	H ₂ O (1:1)/196°C
205	B8a	H	H	Cl	3-(CH ₃ -SO ₂)-phenyl	185°C
206	B8a	CH ₃	H	Cl	phenyl	180°C
207	B8a	H	H	Cl	3-CH ₃ -2-furanyl	188°C
208	B30b	H	H	Cl	3-(CH ₃ -SO ₂ -NH)-phenyl	>250°C
209	B8a	H	H	Cl	2-(CH ₃ -SO ₂)-	230°C
210	B8a	H	H	Cl	2-nitrophenyl	180°C
211	B30a	H	H	Cl	2-aminophenyl	
212	B30b	H	H	Cl	2-(CH ₃ -CO-O-CH ₂ -CO-NH)-phenyl	
213	B26	H	H	Cl	2-(HO-CH ₂ -CO-NH)-phenyl	235°C
214	B30b	H	H	Cl	3-(CH ₃ -CO-O-CH ₂ -CO-NH)-phenyl	
215	B26	H	H	Cl	3-(HO-CH ₂ -CO-NH)-phenyl	>250°C
216	B8a	H	H	Cl	2-CH ₃ -3-nitrophenyl	
217	B30a	H	H	Cl	3-amino-2-methylphenyl	
218	B30b	H	H	Cl	2-CH ₃ -3-(NH ₂ -SO ₂ -NH)-phenyl	180°C
219	B30b	H	H	Cl	3-(C ₂ H ₅ -O-CO-CO-NH)-phenyl	H ₂ O (1:1)/208°C
220	B29b	H	H	Cl		180°C
221	B30b	H	H	Cl	3-(NH ₂ -SO ₂ -NH)-phenyl	H ₂ O (1:1)/220°C
222	B8a	H	H	Cl	2-CH ₃ -3-pyridinyl	160°C
223	B30b	H	H	Cl	2-CH ₃ -3-(CH ₃ -CO-O-CH ₂ -CO-NH)-phenyl	206°C



-54-







Table 3










Co. No.	Ex. No.	R ¹	R ^{4a}	R ^{5a}	R ^{5b}	R ^{11a}	R ^{11b}	Salt form stereochem /mp.
1	B16	CH ₃ O	H	Cl	H	phenyl	H	126°C
2	B14	H	H	Cl	H	H	H	
3	B14	CH ₃	H	Cl	H	H	H	
4	B13a	H	H	Cl	H	phenyl	H	
5	B13a	H	H	Cl	H	4-pyridinyl	H	HBr (1:1)/ H ₂ O (1:1)
6	B13a	H	H	Cl	2-Cl	phenyl	phenyl	
7	B13a	H	H	Cl	2-Cl	phenyl	CH ₃	
8	B13a	CH ₃	H	Cl	2-Cl	phenyl	H	110°C
9	B13a	H	H	Cl	2-Cl	4-Cl-phenyl	H	
10	B13a	H	H	Cl	H	CH ₃	H	
11	B13a	H	H	Cl	H	phenyl	phenyl	
12	B13a	CH ₃	H	Cl	H	phenyl	H	
13	B13a	H	H	Cl	H	4-Cl-phenyl	H	
14	B13a	H	H	Cl	2-Cl	CH ₃	H	
15	B13a	H	H	Cl	2-Cl	4-pyridinyl	H	
16	B13a	H	H	Cl	2-Cl	CH ₃	CH ₃	
17	B13a	H	H	Cl	2-Cl	4-[N(C ₂ H ₅)]-phenyl	H	
18	B13a	CH ₃	H	Cl	2-Cl	phenyl	phenyl	
19	B13a	H	H	Cl	2-Cl	3-Cl-phenyl	H	148°C
20	B13a	H	H	Cl	2-Cl	3-CF ₃ -phenyl	H	155°C
21	B13a	H	H	Cl	2-Cl	3-F-phenyl	H	167°C
22	B13a	H	H	Cl	2-Cl	3-CH ₃ -phenyl	H	162°C
23	B13a	CH ₃	CF ₃	Cl	2-Cl	phenyl	H	130°C
24	B13a	H	H	Cl	2-Cl	3-OCH ₃ -phenyl	H	130°C
25	B13a	H	H	Cl	2-Cl	2-Br-5-OCH ₃ -phenyl	H	130°C
26	B13a	H	H	Cl	2-Cl	4-OH-phenyl	H	255°C
27	B13a	H	H	Cl	2-Cl	C ₂ H ₅ O-CO-	H	220°C
28	B13a	H	H	Cl	2-Cl	3,4-diCl-phenyl	H	170°C
29	B13a	H	H	Cl	2-Cl	phenyl	C ₂ H ₅ O-CO-	144°C

Co. No.	Ex. No.	R ¹	R ^{4a}	R ^{5a}	R ^{5b}	R ^{11a}	R ^{11b}	Salt form stereochem /mp.
30	B13a	H	H	Cl	2-Cl	4-phenyl-phenyl	H	205°C
31	B13a	H	H	Cl	2-Cl	2-thienyl	H	164°C
32	B13a	H	H	Cl	2-Cl	2-Cl-phenyl	H	110°C
33	B4a	OH	H	Cl	H	H	H	
34	B4a	OH	H	Cl	H	phenyl	H	141°C
35	B17b	NH ₂	H	Cl	H	H	H	
36	B17b	Cl	H	Cl	H	H	H	
37	B16	CH ₃ O	H	Cl	H	H	H	
38	B13a	H	H	Cl	2-Cl	phenyl	H	
39	B14	H	H	Cl	2-Cl	H	H	
224	B13a	H	H	Cl	2-Cl	phenyl	ethyl	260°C
225	B13a	H	H	Cl	2-Cl	phenyl-CH ₂ -	H	135°C
226	B13a	H	CF ₃	Cl	2-Cl	phenyl	H	175°C
227	B13a	CH ₃	Cl	Cl	2-Cl	phenyl	H	120°C
228	B13a	CH ₃	Cl	Cl	2-Cl	phenyl	phenyl	130°C
229	B13a	H	H	Cl	2-Cl	4-piperidinyl	H	HCl (1:1)/ 200-210°C
230	B25	H	H	Cl	2-Cl		H	212°C
231	B13a	H	H	Cl	2-Cl	Cl-CH ₂ -	H	
232	B24c	H	H	Cl	2-Cl		H	175°C
233	B13a	CH ₃	Cl	Cl	2-Cl	phenyl	CH ₃	130°C
234	B13a	CH ₃	CF ₃	Cl	2-Cl	phenyl	CH ₃	110°C
235	B13a	CH ₃	CF ₃	Cl	3-CH ₃	phenyl	H	188°C
236	B13a	H	H	Cl	2-Cl	2-furanyl	H	126°C
237	B13a	CH ₃	CF ₃	Cl	2-Cl	phenyl	phenyl	120°C
238	B13a	CH ₃	CF ₃	Cl	3-CH ₃	phenyl	CH ₃	130°C
239	B13a	CH ₃	CF ₃	Cl	H	phenyl	H	126°C
240	B24c	H	H	Cl	2-Cl	(CH ₃) ₂ N-CH ₂ -	H	226°C
241	B13b	H	H	Cl	2-Cl	phenyl	(CH ₃) ₂ CH-	250°C
242	B13a	H	H	Cl	2-Cl	2-F-phenyl	H	85°C
243	B13a	H	H	Cl	2-Cl	2-CH ₃ -phenyl	H	92°C
244	B13a	H	H	Cl	2-Cl	2-Br-phenyl	H	90°C
245	B13a	H	H	Cl	2-Cl	phenyl	propyl	246°C
246	B13a	H	CF ₃	Cl	2-Cl	phenyl	CH ₃	180°C
247	B13a	CH ₃	CH ₃	Cl	H	phenyl	H	150°C


Co. No.	Ex. No.	R ¹	R ^{4a}	R ^{5a}	R ^{5b}	R ^{11a}	R ^{11b}	Salt form stereochem /mp.
248	B13a	H	H	Cl	2-Cl	CH ₃	phenyl	146°C
249	B13a	H	H	Cl	2-Cl	phenyl	phenyl-CH ₂ -	176°C
250	B13a	H	H	Cl	2-Cl	3-Br-phenyl	H	116°C
251	B13a	CH ₃	Cl	Cl	2-Cl	phenyl	ethyl	132°C
252	B13a	H	H	Cl	2-Cl	2,3-diCl-phenyl	H	98°C
253	B13a	H	H	Cl	2-Cl	phenyl	(CH ₃) ₂ N-CH ₂ -	228°C
254	B13a	H	CF ₃	Cl	2-Cl	2-Cl-phenyl	H	104°C
255	B13a	CH ₃	CF ₃	H	2-OCH ₃	phenyl	H	89°C
256	B13a	H	H	Cl	2-Cl	phenyl	C ₂ H ₅ O-CO-CH ₂ -	170°C
257	B13a	H	H	Cl	2-Cl	2,5-diCl-phenyl	H	130°C
258	B13a	H	H	Cl	2-Cl	3-F-phenyl	CH ₃	202°C
259	B13a	H	H	Cl	2-Cl	2-F-phenyl	CH ₃	178°C
260	B13a	H	H	Cl	2-Cl	3-F-phenyl	ethyl	255°C
261	B13a	H	H	Cl	2-Cl	2-F-phenyl	ethyl	152°C
262	B13a	H	H	Cl	Cl	2-Cl-phenyl	ethyl	180°C
263	B13a	H	H	Cl	2-Cl	2-CH ₃ O-phenyl	H	120°C
264	B13a	H	H	Cl	2-Cl	2,6-diCl-phenyl	H	200°C
265	B17a	Cl	H	Cl	H	phenyl	Cl	
266	B3f	(CH ₃) ₂ N- (CH ₂) ₂ -NH-	H	Cl	H	phenyl	Cl	168°C
267	B13a	H	H	Cl	2-Cl	H	phenyl	175°C
268	B13a	H	H	Cl	2-Cl	2,6-diF-phenyl	CH ₃	170°C
269	B13a	H	CH ₃	Cl	2-Cl	phenyl	H	126°C
270	B13a	H	Cl	CH ₃	2-CH ₃	phenyl	H	181°C
271	B13a	H	Cl	CH ₃	2-CH ₃	phenyl	CH ₃	140°C
272	B13a	H	H	Cl	2-Cl	2-Cl-phenyl	CH ₃	182°C
273	B13a	H	H	Cl	2-Cl	phenyl	phenyl-CO-	148°C
274	B13a	H	H	Cl	2-Cl	2-Cl-phenyl	C ₂ H ₅ O-CO-	232°C
275	B13a	H	H	Cl	2-Cl	phenyl	(CH ₃) ₂ N-CO-CH ₂ -	216°C
276	B13a	H	H	Cl	2-Cl	phenyl		203°C
277	B13a	H	H	Cl	2-Cl	phenyl	C ₂ H ₅ O-CO-(CH ₂) ₂ -	184°C
278	B13c	H	H	Cl	2-Cl	phenyl	CH ₃ O-CH ₂ -	228°C
279	B13a	H	H	Cl	2-Cl	phenyl	(CH ₃) ₂ N-(CH ₂) ₂ -	229°C
280	B13a	H	H	Cl	2-Cl	3-F-phenyl	(CH ₃) ₂ N-CH ₂ -	219°C
281	B13a	H	H	Cl	2-Cl	phenyl	(CH ₃) ₂ N-CO-(CH ₂) ₂ -	204°C
282	B24a	H	H	Cl	2-Cl	phenyl	HO-CH ₂ -	142°C
283	B13a	H	H	Cl	2-Cl	phenyl		160°C

Co. No.	Ex. No.	R ¹	R ^{4a}	R ^{5a}	R ^{5b}	R ^{11a}	R ^{11b}	Salt form stereochem /mp.
284	B13a	H	H	Cl	2-Cl	phenyl	cyclohexyl	250°C
285	B13a	H	H	Cl	2-Cl	2-F-phenyl	C ₂ H ₅ O-CO-	222°C
286	B13a	H	H	Cl	2-Cl	3,5-diF-phenyl	H	125°C
287	B13a	H	H	Cl	2-Cl	3-F-phenyl	CH ₃	95°C
288	B13a	CH ₃	F	Cl	2-Cl	phenyl	H	100°C
289	B13a	H	OCH ₃	Cl	2-Cl	phenyl	H	158°C
290	B13a	H	H	Cl	2-Cl	2,5-diF-phenyl	H	120°C
291	B24b	H	H	Cl	2-Cl	phenyl	Cl-CH ₂ -	
292	B24c	H	H	Cl	2-Cl	phenyl		105°C
293	B13a	H	H	Cl	2-Cl	2-Cl-phenyl	C ₂ H ₅ O-CO-CH ₂ -	174°C
294	B13a	H	H	Cl	2-Cl	4-Br-phenyl	H	
295	B13c	H	H	Cl	2-Cl	phenyl	C ₂ H ₅ -O-CH ₂ -	210°C
296	B24c	H	H	Cl	2-Cl	phenyl	CH ₃ -NH-CH ₂ -	HCl (1:1); H ₂ O (1:3)/ 205°C
297	B13a	H	H	Cl	2-Cl	phenyl	phenyl-CH ₂ -N(CH ₃)-CH ₂ -	210°C
298	B13c	H	H	Cl	2-Cl	phenyl	(CH ₃) ₃ C-O-CO-	
299	B22a	H	H	Cl	2-Cl	phenyl	HOOC-	206°C
300	B13a	H	H	Cl	2-Cl	phenyl	HOOC-CH ₂ -	186°C
301	B13c	H	H	Cl	2-Cl	phenyl	CH ₃ -NH-CO-CH ₂ -	158°C
302	B24c	H	H	Cl	2-Cl	phenyl		186°C
303	B22b	H	H	Cl	2-Cl	phenyl	(CH ₃) ₂ N-CO-	150°C
304	B22b	H	H	Cl	2-Cl	phenyl		170°C
305	B22b	H	H	Cl	2-Cl	phenyl		210°C
306	B22b	H	H	Cl	2-Cl	phenyl		156°C
307	B22b	H	H	Cl	2-Cl	phenyl	CH ₃ O-(CH ₂) ₂ -NH-CO-	248°C
308	B13a	H	H	Cl	2-Cl	phenyl	Cl-(CH ₂) ₂ -	
309	B24c	H	H	Cl	2-Cl	phenyl		trifluoroacetate (1:1)
310	B13c	H	H	Cl	2-Cl	phenyl	c.C ₆ H ₁₁ -O-CH ₂ -	200°C
311	B24c	H	H	Cl	2-Cl	phenyl	(CH ₃) ₂ N-(CH ₂) ₂ - N(CH ₃) ₂ -CH ₂ -	170°C

Co. No.	Ex. No.	R ¹	R ^{4a}	R ^{5a}	R ^{5b}	R ^{11a}	R ^{11b}	Salt form stereochem /mp.
312	B22b	H	H	Cl	2-Cl	phenyl	(CH ₃) ₂ N-(CH ₂) ₂ -NH-CO-	H ₂ O (1:1)/ 160°C
313	B24c	H	H	Cl	2-Cl	phenyl		H ₂ O (1:1)/ 216°C
314	B24c	H	H	Cl	2-Cl	phenyl		HCl (1:1)/H ₂ O (1:1)/190°C
315	B24c	H	H	Cl	2-Cl	phenyl	CH ₃ O-CH(CH ₃)-	>260°C
316	B24c	H	H	Cl	2-Cl	phenyl	CH ₃ O-(CH ₂) ₂ -NH-CH ₂ -	110°C
317	B22b	H	H	Cl	2-Cl	phenyl	(CH ₃) ₂ N-(CH ₂) ₂ -NH-CO-CH ₂ -	156°C
318	B13a	H	H	Cl	2-Cl	3-F-phenyl	H	(A)/120°C
319	B13a	H	H	Cl	2-Cl	3-F-phenyl	H	(B)/120°C
320	B22b	H	H	Cl	2-Cl	phenyl	CH ₃ O-(CH ₂) ₂ -NH-CO-CH ₂ -	170-172°C
321	B24c	H	H	Cl	2-Cl	phenyl		210°C
322	B24c	H	H	Cl	2-Cl	phenyl		168°C
323	B24a	H	H	Cl	2-Cl	2-F-phenyl	HO-CH ₂ -	
324	B24b	H	H	Cl	2-Cl	2-F-phenyl	Cl-CH ₂ -	
325	B24c	H	H	Cl	2-Cl	2-F-phenyl	(CH ₃) ₂ N-CH ₂ -	250°C
326	B22b	H	H	Cl	2-Cl	phenyl		140°C
327	B24c	H	H	Cl	2-Cl	phenyl		170°C
328	B13a	H	H	Cl	2-Cl	phenyl		
329	B29a	H	H	Cl	2-Cl	phenyl	NH ₂ -CH ₂ -	H ₂ O (1:1)/ 170°C
330	B13a	H	H	Br	2-Br	phenyl	CH ₃	228°C
331	B29b	H	H	Cl	2-Cl	phenyl	phenyl-NH-C(=S)-NH-CH ₂ -	159°C
332	B24c	H	H	Cl	2-Cl	phenyl	phenyl-(CH ₂) ₂ - N(CH ₃)-CH ₂ -	187°C
333	B29b	H	H	Cl	2-Cl	phenyl	(4-Cl-phenyl)-NH-CO-NH-CH ₂ -	202°C

Co. No.	Ex. No.	R ¹	R ^{4a}	R ^{5a}	R ^{5b}	R ^{11a}	R ^{11b}	Salt form stereochem /mp.
334	B24c	H	H	Cl	2-Cl	phenyl	c.C ₆ H ₁₁ -N(CH ₃)-CH ₂ -	176°C
335	B13a	H	H	Cl	2-Cl	phenyl	(CH ₃) ₂ N-(CH ₂) ₂ -	132°C
336	B30b	H	H	Cl	2-Cl	phenyl	N(CH ₃)-CO-CH ₂ -	158°C
337	B13a	H	H	Cl	2-Cl	2,3-diF-phenyl	phenyl-CH ₂ -SO ₂ -NH-CH ₂ -	110°C


Table 4



Co. No.	Ex. No.	R ^{4a}	R ^{5a}	R ^{11a}	R ^{11b}	R ^{11c}	Salt form / stereochemistry
338	B2a	H	H	OH	c.C ₃ H ₅ -CH ₂ -	CH ₃	CH ₃ SO ₃ H (1:1) H ₂ O (1:2) (A); $\alpha_{20}^D = -346.46^\circ$ (c = 6.35 mg/5 ml in CH ₃ OH) (B); $\alpha_{20}^D = +326.15^\circ$ (c = 6.73 mg/5 ml in CH ₃ OH)
339	B2a	H	H	H	C ₂ H ₅ O-CO-	OH	
340	B2a	H	Cl	H	H	H	
341	B2a	CF ₃	Cl	H	H	H	
342	B2a	CF ₃	Cl	phenyl	H	H	
343	B2a	H	H	H	H	NH ₂	
344	B18	H	Cl	H	H	4-morpholinyl	
345	B18	H	Cl	H	H	4-CH ₃ -1-piperazinyl	
346	B8b	H	Cl	H	H	H	
347	B8b	H	Cl	H	H	H	
348	B18	H	Cl	NH ₂	H	H	
349	B23	H	Cl	H	H	4-morpholinyl	
350	B18	H	Cl	H	H	OH	
351	B18	H	Cl	H	H		

Co. No.	Ex. No.	R ^{4a}	R ^{5a}	R ^{11a}	R ^{11b}	R ^{11c}	Salt form / stereochemistry
352	B18	H	Cl	H	H		HCl (1:1); H ₂ O (1:1)
353	B18	H	Cl	H	H		
354	B18	H	Cl	H	H		
355	B18	H	Cl	(CH ₃) ₂ N-	H	H	
356	B23a	H	Cl	H	H	CF ₃ -SO ₂ -O-	
357	B23b	H	Cl	H	H	HO-(CH ₂) ₂ -NH-	CH ₃ SO ₃ H (1:1)
358	B23b	H	Cl	H	H	[HO-(CH ₂) ₂] ₂ N-	
359	B18	H	Cl		H	H	
360	B27	H	Cl	H	H	1-piperazinyl	H ₂ O (1:1)
361	B23	H	Cl	H	H	(HO-CH ₂) ₂ CH-NH-	
362	B18	H	Cl	H	H		


Table 5



Co. No.	Ex. No.	R ^{5a}	R ^{11a}	R ^{11b}	Salt form / stereochemistry
363	B21	Cl	phenyl	H	HCl (1:1); H ₂ O (1:1) HCl (1:1); H ₂ O (1:1); (A) HCl (1:1); H ₂ O (1:1); (B)
364	B21	Cl	2-F-phenyl	H	
365	B21	Cl	phenyl	CH ₃ -	
366	B21	Cl	4-pyridinyl	H	
366a	B8a	Cl	4-pyridinyl	H	
366b	B8a	Cl	4-pyridinyl	H	
367	B21	Cl	2-Cl-phenyl	H	
368	B21	Cl	3-F-phenyl	H	
369	B21	H	CH ₃	phenyl	
370	B21	Cl	3-F-phenyl	CH ₃ -	
371	B21	Cl	3-Cl-phenyl	H	

Co. No.	Ex. No.	R ^{5a}	R ^{11a}	R ^{11b}	Salt form / stereochemistry
372	B21	Cl	3-CH ₃ -phenyl	H	
373	B21	H	phenyl	phenyl	
374	B21	Cl	2-CH ₃ -phenyl	H	
375	B21	Cl	3-pyridinyl	H	

Table 6




Co. No.	Ex. No.	X	R ²	R ^{4a}	R ^{5a}	salt form / stereochemistry melting point
52	B2a	S	1H-benzimidazol-2-yl	H	H	H ₂ O (1:1)
53	B2a	S	4-CH ₃ -1,2,4-triazol-3-yl	H	H	
54	B2a	S	(CH ₃) ₂ N-(CH ₂) ₂ -	H	H	
55	B2a	S	1H-1,2,4-triazol-3-yl	H	H	
56	B2a	S	5-CH ₃ -1,3,4-thiadiazol-2-yl	H	H	
57	B2a	S	4-F-phenyl	H	H	
58	B2a	S	1-CH ₃ -2-imidazolyl	H	H	
59	B2a	S	4-aminophenyl	H	H	
60	B2a	S	4-OH-6-CH ₃ -2-pyrimidinyl	H	H	
61	B2a	S	4-OH-2-pyrimidinyl	H	H	
62	B2a	S	5-CH ₃ -1H-benzimidazol-2-yl	H	H	
63	B2a	S	2-thiazolyl	H	H	
64	B2a	S	2-furanyl-CH ₂ -	H	H	
65	B2a	S	4-pyridinyl	H	H	
66	B2a	S	4,6-diCH ₃ -2-pyrimidinyl	H	H	
67	B2a	S	4-Cl-phenyl-CH ₂ -	H	H	
68	B2a	S	2,4-diamino-6-pyrimidinyl	H	H	
69	B2a	S	1H-purin-6-yl	H	H	
70	B2a	S	4,6-diamino-2-pyrimidinyl	H	H	
71	B2a	S	2-benzoxazolyl	H	H	
72	B2a	S	4-OH-6-propyl-2-pyrimidinyl	H	H	
73	B2a	S	2-pyridinyl, N-oxide	H	H	
74	B2a	S	1H-pyrazolo[3,4-d]pyrimidin-4-yl	H	H	
75	B2a	S	4-CH ₃ -2-pyrimidinyl	H	H	

Co. No.	Ex. No.	X	R ²	R ^{4a}	R ^{5a}	salt form / stereochemistry melting point
76	B2a	S	C ₂ H ₅ -O-C(=O)-CH ₂ -	H	H	
77	B2a	S	2-benzothiazolyl	H	H	
78	B2a	S	4,5-dihydro-2-thiazolyl	H	H	
79	B2a	S	4-(4-OCH ₃ -phenyl)-2-pyrimidinyl	H	H	
80	B2a	S	CH ₃ -O-C(=O)-(CH ₂) ₂ -	H	H	
81	B2a	S	thiazolo[5,4-b]pyridin-2-yl	H	H	
82	B2a	S	4-OH-6-(CH ₃ OCH ₂)-2-pyrimidinyl	H	H	
83	B2a	S	2-amino-1H-purin-4-yl	H	H	
84	B2a	S	4-(2-thienyl)-2-pyrimidinyl	H	H	
85	B2a	S	6-CH ₃ -5-oxo-4H-1,2,4-triazin-3-yl	H	H	
86	B2a	S	2-pyridinyl	CF ₃	H	
87	B2a	S	4-amino-6-OH-2-pyrimidinyl	H	H	
88	B2a	S	5-CF ₃ -2-pyridinyl	H	H	
89	B2a	S	5-CF ₃ -4H-1,2,4-triazol-3-yl	H	H	
90	B2a	S	cyclohexyl	H	H	
91	B2a	S	5-ethyl-4-oxo-2(3H)-pyrimidinyl	H	H	
92	B1b	S	2-pyrimidinyl	H	H	
93	B2a	S	2-pyridinyl	H	H	
94	B2b	S	1H-imidazol-2-yl	H	H	
95	B2c	S	C ₂ H ₅ -O-C(=O)-CH(NH ₂)-	H	H	
96	B11	S	2,4-diOCH ₃ -6-pyrimidinyl	H	H	
98	B1	O	CH ₃	H	H	
133	B1	O	(CH ₃) ₂ CH-CH ₂	H	H	
376	B2a	S	thiazolo[5,4-b]pyridin-2-yl	H	Cl	
377	B2a	S	2-pyridinyl	H	Cl	
377a	B8a	S	2-pyridinyl	H	Cl	(A); $\alpha_{20}^D = +354.70^\circ$ (c = 5.85 mg/5 ml in CH ₃ OH)
377b	B8a	S	2-pyridinyl	H	Cl	(B); $\alpha_{20}^D = -356.73^\circ$ (c = 6.91 mg/5 ml in CH ₃ OH)
378	B2a	S	2-pyridinyl	CF ₃	Cl	
379	B2a	S	2-benzoxazolyl	CF ₃	Cl	
380	B2a	S	4-phenyl-2-thiazolyl	H	Cl	
381	B2a	S	4-phenyl-2-thiazolyl	CF ₃	Cl	
382	B2a	S	thiazolo[5,4-b]pyridin-2-yl	CF ₃	Cl	
383	B2a	S	2-benzoxazolyl	H	Cl	
384	B2a	S	2-benzothiazolyl	H	Cl	
385	B2a	S	2-benzothiazolyl	CF ₃	Cl	

Co. No.	Ex. No.	X	R ²	R ^{4a}	R ^{5a}	salt form / stereochemistry melting point
386	B2a	S	4,5-dihydro-2-thiazolyl	CF ₃	Cl	216°C
387	B2a	S	2-thiazolyl	CF ₃	Cl	
388	B2a	S	6-nitro-2-benzothiazolyl	CF ₃	Cl	
389	B2a	S	6-NH ₂ -2-benzothiazolyl	CF ₃	Cl	
390	B2a	S	4-(2-thienyl)-2-thiazolyl	CF ₃	Cl	
391	B2a	S	5-phenyl-1,3,4-oxadiazol-2-yl	CF ₃	Cl	
392	B2a	S	5CH ₃ -4-phenyl-2-thiazolyl	CF ₃	Cl	
393	B2a	S	4-NH ₂ -phenyl	CF ₃	Cl	
394	B2a	S	6-ethoxy-2-benzothiazolyl	CF ₃	Cl	
395	B2a	S	pyrido[3,4-d]thiazol-2-yl	CF ₃	Cl	
396	B2a	S	1H-benzimidazol-2-yl	CF ₃	Cl	
397	B2a	S	4-(2,4-diF-phenyl)-2-thiazolyl	CF ₃	Cl	
398	B2a	S	4-(CH ₃ -CO-NH)-phenyl	CF ₃	Cl	
399	B2a	S	4-(2-furanyl)-2-thiazolyl	CF ₃	Cl	
400	B2d	S	1,3-dihydro-4-phenyl-2H-imidazole-2-thion-5-yl	CF ₃	Cl	
401	B2a	S	2-pyrazinyl	CF ₃	Cl	
402	B2a	S	5-Cl-2-benzothiazolyl	CF ₃	Cl	
403	B2a	S	pyrido[3,4-d]oxazol-2-yl	CF ₃	Cl	
404	B2a	S	3-phenyl-1,2,4-oxadiazol-5-yl	CF ₃	Cl	
405	B2a	S	5-CH ₃ -4-phenyl-2-thiazolyl	CF ₃	Cl	
406	B18	S	5-phenyl-1,3,4-oxadiazol-2-yl	H	Cl	
407	B2a	S	(2-pyrazinyl)-CH ₂ -	H	Cl	
408	B18	S	3-phenyl-1,2,4-oxadiazol-5-yl	H	Cl	
409	B18	S	4-pyrimidinyl	H	Cl	


Table 7



Co. No.	Ex. No.	R ²	R ^{4a}	R ^{5a}	salt form
40	B3e	5-CH ₃ -3-isoxazolyl	H	H	HCl (1:1)
41	B3c	CH ₃ -O-(CH ₂) ₂ -	H	H	
42	B3c	4-CH ₃ -6-OCH ₃ -2-pyrimidinyl	H	H	
43	B3c	2-furanylethyl	H	H	
44	B3c	2-thiazolyl	H	H	

Co. No.	Ex. No.	R ²	R ^{4a}	R ^{5a}	salt form
46	B3a	cyclohexyl	H	H	
47	B10b	benzoyl	H	H	
48	B3f	1-CH ₃ -4-piperidiny	H	H	
49	B3e	2-pyrimidinyl	H	H	
50	B3d	1 <i>H</i> -imidazol-2-yl	H	H	
51	B3c	C ₂ H ₄ OH	H	H	
410	B10b	thiazolo[5,4-b]pyridin-2-yl	H	H	
411	B3g	4-phenyl-2-thiazolyl	CF ₃	Cl	
412	B3c	5-CH ₃ -4-phenyl-2-thiazolyl	H	H	
413	B3g	2-pyrimidinyl	H	Cl	

Table 8



Co. No.	Ex. No.	R ¹	R ²	R ^{4a}	R ^{5a}	R ^{5b}	Salt form melting point
45	B3a	H	N(CH ₃) ₂	H	Cl	H	
97	B3c	H	1,2,4-triazol-1-yl	H	Cl	H	
99	B3c	H	1,2,4-triazol-4-yl	H	Cl	H	
100	B3c	H	1 <i>H</i> -imidazol-1-yl	H	Cl	H	
101	B8a	H	5-phenyl-1,3,4-oxadiazol-2-yl	H	Cl	H	
102	B8a	H	5-CH ₃ -1,3,4-oxadiazol-2-yl	H	Cl	H	
103	B8a	H	5-phenyl-2-oxazolyl	H	Cl	H	
104	B8a	CH ₃	5-phenyl-1,3,4-oxadiazol-2-yl	H	Cl	H	
105	B8a	H	5-phenyl-2-oxazolyl	H	Cl	Cl	
106	B6	CH ₃	3-phenyl-1,2,4-oxadiazol-5-yl	H	Cl	H	
107	B7	H	5-phenyl-1,2,4-oxadiazol-3-yl	H	Cl	H	
108	B5a	H	2-CH ₃ -1,2,4-triazol-3-yl	H	Cl	H	
109	B5a	H	1-CH ₃ -2-imidazolyl	H	Cl	Cl	164°C
110	B4a	OH	2-CH ₃ -1,2,4-triazol-3-yl	H	Cl	H	H ₂ O (1:1)
111	B4a	OH	2-benzothiazolyl	H	Cl	H	
112	B5a	H	4-pyridinyl	H	Cl	H	
113	B5a	H	4-pyridinyl	H	Cl	Cl	
114	B5a	H	2-pyridinyl	H	Cl	H	130°C
115	B5a	H	2-pyridinyl	H	Cl	Cl	205°C
116	B5a	H	3-pyridinyl	H	Cl	Cl	166°C
117	B4a	OH	3-pyridinyl	H	Cl	H	

Co. No.	Ex. No.	R ¹	R ²	R ^{4a}	R ^{5a}	R ^{5b}	Salt form melting point
118	B3a	H	4-CH ₃ -1-piperazinyl	H	Cl	H	H ₂ O (1:1) H ₂ O (1:1)
119	B3b	H	4-OH-1-piperidinyl	H	Cl	H	
120	B4a	OH	1-CH ₃ -2-imidazolyl	H	Cl	H	
121	B4b	OH	3-CH ₃ -4-imidazolyl	H	Cl	H	
122	B4c	OH	CN-CH ₂ -	H	Cl	Cl	
123	B5a	H	1-CH ₃ --2-imidazolyl	H	Cl	H	
124	B5b	H	3-pyridinyl	H	Cl	H	
125	B6	H	3-phenyl-1,2,4-oxadiazol-5-yl	H	Cl	H	
126	B7	H	5-CH ₃ -1,2,4-oxadiazol-3-yl	H	Cl	H	
127	B8a	H	5-phenyl-1,3,4-oxadiazol-2-yl	H	Cl	Cl	
128	B9	H	5-SH-4-phenyl-1,2,4-triazol-3-yl	H	Cl	H	240°C 128°C 140°C 100°C 90°C 208°C
129	B9	H	5-(phenyl-NH)-1,3,4-thiadiazol-2-yl	H	Cl	H	
130	B12	H	2-benzothiazolyl	H	Cl	H	
131	B15a	H	2-benzoxazolyl	H	Cl	H	
132	B15b	CH ₃	2-benzoxazolyl	H	Cl	H	
414	B12	H	5-phenyl-1,3,4-thiadiazol-2-yl	H	H	Cl	
415	B17a	Cl	2-benzothiazolyl	H	Cl	H	
416	B17b	NH ₂	2-benzothiazolyl	H	Cl	H	
417	B4c	HO	CN-CH ₂ -	CF ₃	Cl	Cl	
418	B16	CH ₃ O	2-benzothiazolyl	H	Cl	H	
419	B20	H	(4-phenyl-2-thiazolyl)-CH ₂ -	H	H	Cl	>250°C 150°C 140°C H ₂ O(1:1)/245°C 128°C >260°C 220°C
420	B19a	H	HO-CH ₂ -	H	Cl	Cl	
421	B5a	H	2-benzothiazolyl	H	Cl	Cl	
422	B19c	H	(2-pyrimidinyl)thio-CH ₂ -	H	H	Cl	
423	B19a	H	HO-CH ₂ -	CF ₃	Cl	Cl	
424	B19b	H	H ₃ C-SO ₂ -O-CH ₂ -	CF ₃	Cl	Cl	
425	B5a	H	1-CH ₃ -4-phenyl-2-imidazolyl	H	Cl	Cl	
426	B8a	H	5-CH ₃ -4-phenyl-2-oxazolyl	H	Cl	Cl	
427	B12	H	5-phenyl-1,3,4-thiadiazol-2-yl	H	Cl	Cl	
428	B5a	H	4-CH ₃ -5-phenyl-1,2,4-triazol-3-yl	H	Cl	Cl	
429	B6b	H	3-phenyl-1,2,4-oxadiazol-5-yl	H	Cl	Cl	195°C
430	B5a	H	1-CH ₃ -2-phenyl-5-imidazolyl	H	Cl	Cl	
431	B8a	H	5-CH ₃ -4-(4-F-phenyl)-2-oxazolyl	H	Cl	Cl	
432	B21	H	5-phenylimidazo[2,1-b]thiazol-6-yl	H	H	Cl	
433	B21	H	5,6-dihydro-2-phenylimidazo-[2,1-b]thiazol-3-yl	H	H	Cl	
434	B5a	H	2,4-diphenyl-5-oxazolyl	H	Cl	Cl	
435	B19b	H	H ₃ C-SO ₂ -O-CH ₂ -	H	Cl	Cl	

C. Pharmacological example

Example C.1 : *in vitro* inhibition of IL-5 production in human blood

Human whole blood stimulation

- Peripheral blood from healthy male donors was drawn into heparinized syringes (12.5 U heparin/ml). Blood samples were three-fold diluted in RPMI 1640 medium (Life Technologies, Belgium) supplemented with 2 mM L-glutamine, 100 U/ml penicillin and 100 µg/ml streptomycin, and 300 µl fractions were distributed in 24-well multidisc plates. Blood samples were preincubated (60 minutes at 37°C) in a humidified 6% CO₂-atmosphere with 100 µl of drug solvent (final concentration 0.02% dimethylsulfoxide in RPMI 1640) or with 100 µl of an appropriate dose of test compound before being stimulated by the addition of 100 µl of phytohemagglutinin HA17 (Murex, UK) at a final concentration of 2 µg/ml. After 48 hours, cell-free supernatant fluids were collected by centrifugation and stored at -70°C until tested for the presence of IL-5.

15 *IL-5 measurements*

IL-5 measurements were conducted as described in Van Wauwe et al. (1996, *Inflamm Res*, 45, 357-363) on page 358 using ELISA.

- Table 9 lists the percentage inhibition of IL-5 production (column "% inh") at a test dose of 1×10^{-6} M, or in case the percentage inhibition is marked with an "*" 1×10^{-5} M, for the compounds of the present invention.

Table 9

Comp No	% inh.	Comp No	% inh.	Comp No	% inh.	Comp No	% inh.	Comp No	% inh.	Comp No	% inh.
1	77	18	88	37	61	55	26	74	69	95	17
2	55*	19	83	38	92	56	41	75	72	96	62
3	46	20	70	39	68	57	50	76	2	97	26
4	83	21	91	40	31	58	5	77	65	101	66
5	77	22	93	41	11	59	76	78	70	102	14
6	91	23	83	42	57	60	24*	79	74	103	63
7	95	24	74	43	37	61	14	81	76	104	60
8	93	25	88	44	40	62	30	82	19	105	88
9	85	26	85	46	64	63	68	84	73	106	77
10	64	27	64	47	33	64	64	85	38	107	81
11	91	28	73	48	29	65	50	86	84	109	35
12	77	29	95	49	61	66	64	87	9	110	6
13	61	30	57	50	20*	67	69	88	26	111	61
14	83	31	93	51	10	68	60	89	19	112	62
15	86	32	90	52	57*	70	51	90	60	113	76
16	89	34	58	53	53*	71	84	93	86	114	40
17	81	35	56	54	14	73	21	94	18	115	71

Comp No	% inh.
116	74
117	34
118	34
119	72*
120	10
123	13
124	42
125	52
126	40
127	94
130	70
131	76
132	55
133	50
134	95
135	88
136	93
137	64
138	81
139	60
140	45
141	64
142	80
143	81
144	40
145	37
146	83
147	50
148	79
149	89
150	48
151	17
152	87
153	72
154	42
155	80
156	91
157	85
158	92
159	87
160	91
161	91
162	63
163	90
164	84
165	80

Comp No	% inh.
166	87
167	82
168	80
169	81
170	62
171	59
172	17
173	44
174	83
175	58
176	3
177	69
178	78
179	21
180	54
181	55
182	75
184	83
185	81
186	8
187	25
188	95
189	82
190	83
191	19
194	83
195	7
196	35
198	46
199	43
200	43
201	87
203	82
204	36
205	80
206	82
207	94
208	48
209	77
210	79
211	83
213	32
215	54
218	4
219	8
220	25

Comp No	% inh.
221	-2
224	95
225	80
226	93
227	78
228	81
230	79
232	47
233	84
234	83
235	79
236	92
237	82
238	74
239	72
240	54
241	95
242	98
243	97
244	95
245	98
246	94
247	80
248	91
249	80
250	84
251	90
252	80
253	96
254	86
255	67
256	94
257	82
258	98
259	95
260	98
261	93
262	93
263	92
264	79
266	46
267	81
268	83
269	90
270	86
271	88

Comp No	% inh.
272	87
273	77
274	89
275	94
276	91
277	66
278	97
279	92
280	96
281	91
282	93
283	93
284	91
285	89
286	86
287	94
288	90
289	96
290	92
292	94
293	59
294	85
295	90
296	92
297	90
299	38
300	27
301	33
302	87
303	85
304	35
305	51
306	92
307	78
309	82
310	79
311	64
312	57
313	86
314	81
315	93
316	85
317	67
318	81
319	84
320	94

Comp No	% inh.
321	92
322	96
325	95
326	89
327	84
329	88
330	94
331	95
332	86
333	61
334	75
335	52
336	88
337	96
338	-15
339	35
340	88
341	96
342	93
343	66
344	82
345	88
346	86
347	8
348	83
349	87
351	62
352	85
353	91
354	70
355	83
357	69
358	63
359	88
360	84
361	28
363	91
364	95
365	88
366	93
367	74
368	88
369	66
370	76
371	88
372	86

Comp No	% inh.
373	40
374	94
375	91
376	92
377	87
378	91
379	95
380	95
381	95
382	95
383	78
384	95
385	95
386	97
387	93
388	90
389	91
390	89
391	97
392	87
393	93
394	93
395	94
396	28
397	83
398	96
399	93
400	76
401	92
402	90
403	97
404	92
405	80
406	84
407	71
408	88
409	88
410	15
411	94
412	16
413	59
414	30
416	79
418	47
419	5
420	33

Comp No	% inh.
421	86
422	87

Comp No	% inh.
425	70
426	92

Comp No	% inh.
427	72
428	66

Comp No	% inh.
429	78
430	89

Comp No	% inh.
431	67
432	82

Comp No	% inh.
433	53
434	72

D. Composition examples

The following formulations exemplify typical pharmaceutical compositions suitable for systemic or topical administration to animal and human subjects in accordance with the present invention.

- 5 "Active ingredient" (A.I.) as used throughout these examples relates to a compound of formula (I) or a pharmaceutically acceptable addition salt thereof.

Example D.1 : film-coated tablets

Preparation of tablet core

- 10 A mixture of A.I. (100 g), lactose (570 g) and starch (200 g) was mixed well and thereafter humidified with a solution of sodium dodecyl sulfate (5 g) and polyvinyl-pyrrolidone (10 g) in about 200 ml of water. The wet powder mixture was sieved, dried and sieved again. Then there was added microcrystalline cellulose (100 g) and hydrogenated vegetable oil (15 g). The whole was mixed well and compressed into tablets, giving 10.000 tablets, each comprising 10 mg of the active ingredient.

Coating


- 15 To a solution of methyl cellulose (10 g) in denaturated ethanol (75 ml) there was added a solution of ethyl cellulose (5 g) in CH_2Cl_2 (150 ml). Then there were added CH_2Cl_2 (75 ml) and 1,2,3-propanetriol (2.5 ml). Polyethylene glycol (10 g) was molten and dissolved in dichloromethane (75 ml). The latter solution was added to the former and then there were
20 added magnesium octadecanoate (2.5 g), polyvinyl-pyrrolidone (5 g) and concentrated color suspension (30 ml) and the whole was homogenated. The tablet cores were coated with the thus obtained mixture in a coating apparatus.

Example D.2 : 2% topical cream

- 25 To a solution of hydroxypropyl β -cyclodextrine (200 mg) in purified water is added A.I. (20 mg) while stirring. Hydrochloric acid is added until complete dissolution and next sodium hydroxide is added until pH 6.0. While stirring, glycerol (50 mg) and polysorbate 60 (35 mg) are added and the mixture is heated to 70°C. The resulting mixture is added to a mixture of mineral oil (100 mg), stearyl alcohol (20 mg), cetyl alcohol (20 mg), glycerol monostearate (20 mg) and sorbate 60 (15 mg) having a
30 temperature of 70°C while mixing slowly. After cooling down to below 25°C, the rest of the purified water q.s. ad 1 g is added and the mixture is mixed to homogenous.

Claims

1. A compound of formula



- a *N*-oxide, a pharmaceutically acceptable addition salt or a stereochemically isomeric form thereof, wherein :
- p* represents an integer being 0, 1, 2, 3 or 4;
- q* represents an integer being 0, 1, 2, 3, 4 or 5;
- X* represents O, S, NR³ or a direct bond;
- R¹ represents hydrogen, hydroxy, halo, amino, mono- or di(C₁₋₄alkyl)amino, C₁₋₆alkyl, C₁₋₆alkyloxy, C₃₋₇cycloalkyl, aryl, arylC₁₋₆alkyl, aminoC₁₋₄alkyl, mono- or di(C₁₋₄alkyl)aminoC₁₋₄alkyl or mono- or di(C₁₋₄alkyl)aminoC₁₋₄alkylamino;
- R² represents aryl, Het¹, C₃₋₇cycloalkyl, C₁₋₆alkyl or C₁₋₆alkyl substituted with one or two substituents selected from hydroxy, cyano, amino, mono- or di(C₁₋₄alkyl)amino, C₁₋₆alkyloxy, C₁₋₆alkylsulfonyloxy, C₁₋₆alkyloxycarbonyl, C₃₋₇cycloalkyl, aryl, aryloxy, arylthio, Het¹, Het¹oxy and Het¹thio; and if *X* is O, S or NR³, then R² may also represent aminocarbonyl, aminothiocarbonyl, C₁₋₄alkylcarbonyl, C₁₋₄alkylthiocarbonyl, arylcarbonyl or arylthiocarbonyl;
- R³ represents hydrogen or C₁₋₄alkyl;
- each R⁴ independently represents C₁₋₆alkyl, halo, polyhaloC₁₋₆alkyl, hydroxy, mercapto, C₁₋₆alkyloxy, C₁₋₆alkylthio, C₁₋₆alkylcarbonyloxy, aryl, cyano, nitro, Het³, R⁶, NR⁷R⁸ or C₁₋₄alkyl substituted with Het³, R⁶ or NR⁷R⁸;
- each R⁵ independently represents C₁₋₆alkyl, halo, polyhaloC₁₋₆alkyl, hydroxy, mercapto, C₁₋₆alkyloxy, C₁₋₆alkylthio, C₁₋₆alkylcarbonyloxy, aryl, cyano, nitro, Het³, R⁶, NR⁷R⁸ or C₁₋₄alkyl substituted with Het³, R⁶ or NR⁷R⁸;
- each R⁶ independently represents C₁₋₆alkylsulfonyl, aminosulfonyl, mono- or di(C₁₋₄alkyl)aminosulfonyl, mono- or di(benzyl)aminosulfonyl, polyhaloC₁₋₆alkylsulfonyl, C₁₋₆alkylsulfinyl, phenylC₁₋₄alkylsulfonyl, piperazinylsulfonyl, aminopiperidinylsulfonyl, piperidinylaminosulfonyl, *N*-C₁₋₄alkyl-*N*-piperidinylaminosulfonyl;
- each R⁷ and each R⁸ are independently selected from hydrogen, C₁₋₄alkyl, hydroxyC₁₋₄alkyl, dihydroxyC₁₋₄alkyl, aryl, arylC₁₋₄alkyl, C₁₋₄alkyloxyC₁₋₄alkyl, C₁₋₄alkylcarbonyl, arylcarbonyl, C₁₋₄alkylcarbonyloxyC₁₋₄alkylcarbonyl, hydroxyC₁₋₄alkylcarbonyl, C₁₋₄alkyloxycarbonylcarbonyl, mono- or

- di(C₁₋₄alkyl)aminoC₁₋₄alkyl, arylaminocarbonyl, arylaminothiocarbonyl, Het³aminocarbonyl, Het³aminothiocarbonyl, C₃₋₇cycloalkyl, pyridinylC₁₋₄alkyl, Het³ and R⁶;
- R⁹ and R¹⁰ are each independently selected from hydrogen, C₁₋₄alkyl, hydroxyC₁₋₄alkyl, dihydroxyC₁₋₄alkyl, phenyl, phenylC₁₋₄alkyl, C₁₋₄alkyloxyC₁₋₄alkyl, C₁₋₄alkylcarbonyl, phenylcarbonyl, C₁₋₄alkylcarbonyloxyC₁₋₄alkylcarbonyl, hydroxyC₁₋₄alkylcarbonyl, C₁₋₄alkyloxycarbonylcarbonyl, mono- or di(C₁₋₄alkyl)aminoC₁₋₄alkyl, phenylaminocarbonyl, phenylaminothiocarbonyl, Het³aminocarbonyl, Het³aminothiocarbonyl, C₃₋₇cycloalkyl, pyridinylC₁₋₄alkyl, Het³ and R⁶;
- each R¹¹ independently being selected from hydroxy, mercapto, cyano, nitro, halo, trihalomethyl, C₁₋₄alkyloxy, carboxyl, C₁₋₄alkyloxycarbonyl, trihaloC₁₋₄alkylsulfonyloxy, R⁶, NR⁷R⁸, C(=O)NR⁷R⁸, aryl, aryloxy, arylcarbonyl, C₃₋₇cycloalkyl, C₃₋₇cycloalkyloxy, phthalimide-2-yl, Het³ and C(=O)Het³;
- R¹² and R¹³ are each independently selected from hydrogen, C₁₋₄alkyl, hydroxyC₁₋₄alkyl, dihydroxyC₁₋₄alkyl, phenyl, phenylC₁₋₄alkyl, C₁₋₄alkyloxyC₁₋₄alkyl, C₁₋₄alkylcarbonyl, phenylcarbonyl, C₁₋₄alkylcarbonyloxyC₁₋₄alkylcarbonyl, hydroxyC₁₋₄alkylcarbonyl, C₁₋₄alkyloxycarbonylcarbonyl, mono- or di(C₁₋₄alkyl)aminoC₁₋₄alkyl, phenylaminocarbonyl, phenylaminothiocarbonyl, C₃₋₇cycloalkyl, pyridinylC₁₋₄alkyl and R⁶;
- aryl represents phenyl optionally substituted with one, two or three substituents each independently selected from nitro, azido, halo, hydroxy, C₁₋₄alkyl, C₁₋₄alkyloxy, polyhaloC₁₋₄alkyl, NR⁹R¹⁰, R⁶, phenyl, Het³ and C₁₋₄alkyl substituted with NR⁹R¹⁰;
- Het¹ represents a heterocycle selected from pyrrolyl, pyrrolinyl, imidazolyl, imidazolinyl, pyrazolyl, pyrazolinyl, triazolyl, tetrazolyl, furanyl, tetrahydrofuranyl, thienyl, thiolanyl, dioxolanyl, oxazolyl, oxazolinyl, isoxazolyl, thiazolyl, thiazolinyl, isothiazolyl, thiadiazolyl, oxadiazolyl, pyridinyl, pyrimidinyl, pyrazinyl, pyranyl, pyridazinyl, pyrrolidinyl, piperidinyl, piperazinyl, morpholinyl, thiomorpholinyl, dioxanyl, dithianyl, trithianyl, triazinyl, benzothienyl, isobenzothienyl, benzofuranyl, isobenzofuranyl, benzothiazolyl, benzoxazolyl, indolyl, isoindolyl, indolinyl, purinyl, 1H-pyrazolo[3,4-d]pyrimidinyl, benzimidazolyl, quinolyl, isoquinolyl, cinnolinyl, phtalazinyl, quinazolinyl, quinoxalinyl, thiazolopyridinyl, oxazolopyridinyl, imidazo[2,1-b]thiazolyl; wherein said heterocycles each independently may optionally be substituted with one, or where possible, two or three substituents each independently selected from Het², R¹¹ and C₁₋₄alkyl optionally substituted with Het² or R¹¹;
- Het² represents a monocyclic heterocycle selected from pyrrolyl, pyrrolinyl, imidazolyl, imidazolinyl, pyrazolyl, pyrazolinyl, triazolyl, tetrazolyl, furanyl, tetrahydrofuranyl, thienyl, thiolanyl, dioxolanyl, oxazolyl, oxazolinyl, isoxazolyl, thiazolyl, thiazolinyl,


isothiazolyl, thiadiazolyl, oxadiazolyl, pyridinyl, pyrimidinyl, pyrazinyl, pyranyl, pyridazinyl, dioxanyl, dithianyl, trithianyl and triazinyl; wherein said monocyclic heterocycles each independently may optionally be substituted with one, or where possible, two or three substituents each independently selected from R^{11} and C_{1-4} alkyl optionally substituted with R^{11} ;

Het³ represents a monocyclic heterocycle selected from pyrrolidinyl, piperidinyl, piperazinyl, morpholinyl, thiomorpholinyl; wherein said monocyclic heterocycles each independently may optionally be substituted with, where possible, one, two or three substituents each independently selected from C_{1-4} alkyl, C_{1-4} alkyloxy, carboxyl, C_{1-4} alkyloxycarbonyl, C_{1-4} alkylcarbonyl, phenyl C_{1-4} alkyl, piperidinyl, $NR^{12}R^{13}$, R^6 and C_{1-4} alkyl substituted with R^6 or $NR^{12}R^{13}$.


2. A compound according to claim 1 wherein R^1 is hydrogen, hydroxy, halo, amino, C_{1-6} alkyl, C_{1-6} alkyloxy or mono- or di(C_{1-4} alkyl)amino C_{1-4} alkylamino.
3. A compound according to claim 1 or 2 wherein R^2 is aryl, Het¹, C_{3-7} cycloalkyl, or C_{1-6} alkyl substituted with one or two substituents selected from hydroxy, cyano, amino, mono- or di(C_{1-4} alkyl)amino, C_{1-6} alkyloxy, C_{1-6} alkylsulfonyloxy, C_{1-6} alkyloxycarbonyl, C_{3-7} cycloalkyl, aryl, aryloxy, arylthio, Het¹, Het¹oxy and Het¹thio; and if X is O, S or NR^3 , then R^2 may also represent aminocarbonyl, aminothiocarbonyl, C_{1-4} alkylcarbonyl, C_{1-4} alkylthiocarbonyl, arylcarbonyl or arylthiocarbonyl.
4. A compound according to any one of claims 1 to 3 wherein the 6-azauracil moiety is in the para position relative to the central carbon atom.
5. A compound according to any one of claims 1 to 4 wherein q is 1 or 2 and one R^4 substituent is in the 4 position; and p is 1 or 2 and the one or two R^5 substituents are in the ortho position relative to the central carbon atom.
6. A composition comprising a pharmaceutically acceptable carrier and, as active ingredient, a therapeutically effective amount of a compound as claimed in any one of claims 1 to 5.
7. A process for preparing a composition as claimed in claim 6, wherein a pharmaceutically acceptable carrier is intimately mixed with a therapeutically effective amount of a compound as defined in any one of claims 1 to 5.
8. A compound as claimed in any one of claims 1 to 5 for use as a medicine.

9. Use of a compound as claimed in any one of claims 1 to 5 in the manufacture of a medicament for treating eosinophil-dependent inflammatory diseases.

- 5 10. A process for preparing a compound as claimed in claim 1, characterized by,
a) reacting an intermediate of formula (II) wherein W^1 is a suitable leaving group with an appropriate reagent of formula (III) optionally in a reaction-inert solvent and in the presence of a base;




- 10 wherein R^1 , R^2 , R^4 , X and q are as defined in claim 1, and D represents




wherein R^5 and p are defined as in claim 1;

- b) eliminating the group E of a triazinedione of formula (V)




- 15 wherein R^1 , R^2 , R^4 , R^5 , X and q are as defined in claim 1;

c) reacting a ketone of formula (X) with an intermediate of formula (III-a) in the presence of a base and in a reaction-inert solvent; thus obtaining a compound of formula (I-a-2);




- 20 wherein R^2 , R^4 and q are as defined in claim 1 and D is defined as in claim 9a);

d) converting a compound of formula (I-a-2) to a compound of formula (I-a-3) using art-known group transformation reactions,




wherein R^2 , R^4 and q are as defined in claim 1 and D is defined as in claim 9a);

5 e) converting a compound of formula (I-a-2) to a compound of formula (I-a-4) using art-known group transformation reactions,




wherein R^2 , R^4 and q are as defined in claim 1 and D is defined as in claim 9a);

10 f) converting a compound of formula (I-a-4) to a compound of formula (I-a-5) using art-known group transformation reactions,



wherein R^2 , R^4 and q are as defined in claim 1 and D is defined as in claim 9a);


g) reacting an intermediate of formula (XII) wherein W^4 is a suitable leaving group with an intermediate of formula (III) optionally in the presence of a suitable base; thus
15 obtaining a compounds of formula (I-b);



wherein R^2 , R^4 , X and q are as defined in claim 1 and D is defined as in claim 9a);


h) reacting an intermediate of formula (XIV) with an intermediate of formula (XV) wherein W^3 is a suitable leaving group, in the presence of a suitable base and

optionally in the presence of a reaction-inert solvent; thus obtaining a compound of formula (I-c);




wherein R^4 and q are as defined in claim 1 and D is defined as in claim 9a);

- 5 i) cyclizing an intermediate of formula (XX) wherein Y is O, S or NR^3 , to a compound of formula (I-d-1), in the presence of a suitable solvent at an elevated temperature;




wherein R , R^1 , R^4 and q are as defined in claim 1 and D is defined as in claim 9a);

- 10 j) cyclizing an intermediate of formula (XXI) to a compound of formula (I-d-2) in a reaction-inert solvent at an elevated temperature,



wherein R , R^1 , R^4 and q are as defined in claim 1 and D is defined as in claim 9a);


- k) cyclizing an intermediate of formula (XXII) wherein Y is O, S or NR^3 , to a compound of formula (I-d-3), in a suitable solvent,



15


wherein R , R^1 , R^4 and q are as defined in claim 1 and D is defined as in claim 9a);

l) cyclizing an intermediate of formula (XXIII) wherein Y is O, S or NR³, to a compound of formula (I-d-4), in a reaction-inert solvent and in the presence of an acid,




wherein R, R¹, R⁴ and q are as defined in claim 1 and D is defined as in claim 9a);

- 5 m) cyclizing an intermediate of formula (XXIII) wherein Y is O, S or NR³, to a compound of formula (I-d-5), in a reaction-inert solvent and in the presence of an acid,



wherein R, R¹, R⁴ and q are as defined in claim 1 and D is defined as in claim 9a);


- n) reacting an intermediate of formula (XXIV) with an intermediate of formula (XXV) wherein Y is O, S or NR³, and W⁵ is a suitable leaving group; thus forming a compound of formula (I-d-6) in a reaction-inert solvent and in the presence of a base,



wherein R, R¹, R⁴ and q are as defined in claim 1 and D is defined as in claim 9a);


- o) reacting an intermediate of formula (XXVI) with an intermediate of formula (XXVII) wherein W⁶ is a suitable leaving group; thus forming a compound of formula (I-d-7), in a reaction-inert solvent and in the presence of an acid;

-76-



wherein R, R¹, R⁴ and q are as defined in claim 1 and D is defined as in claim 9a);

- p) reacting an intermediate of formula (XXXIII) with a thioamide of formula (XXXIV); thus forming a compound of formula (I-d-9) in a reaction-inert solvent at an elevated temperature;



wherein R, R¹, R⁴ and q are as defined in claim 1 and D is defined as in claim 9a);

- and if desired, converting compounds of formula (I) into each other following art-known transformations, and further, if desired, converting the compounds of formula (I), into a therapeutically active non-toxic acid addition salt by treatment with an acid, or into a therapeutically active non-toxic base addition salt by treatment with a base, or conversely, converting the acid addition salt form into the free base by treatment with alkali, or converting the base addition salt into the free acid by treatment with acid; and also, if desired, preparing stereochemically isomeric forms or *N*-oxide forms thereof.

15

11. A process of marking a receptor comprising the steps of
 - a) radiolabelling a compound as defined in claim 1;
 - b) administering said radiolabelled compound to biological material,
 - c) detecting the emissions from the radiolabelled compound.

20

12. A process of imaging an organ, characterized by, administering a sufficient amount of a radiolabelled compound of formula (I) in an appropriate composition, and detecting the emissions from the radioactive compound.

INTERNATIONAL SEARCH REPORT

International Application No

PCT/EP 98/04191

A. CLASSIFICATION OF SUBJECT MATTER

IPC 6 C07D253/075 C07D401/12 C07D403/12 C07D413/10 C07D417/10
 C07D495/04 C07D409/10 C07D413/12 C07D417/14 A61K31/53
 C07D401/14 C07D413/14 //(C07D495/04,213:00,333:00)

According to International Patent Classification (IPC) or to both national classification and IPC

B. FIELDS SEARCHED

Minimum documentation searched (classification system followed by classification symbols)

IPC 6 C07D A61K

Documentation searched other than minimum documentation to the extent that such documents are included in the fields searched

Electronic data base consulted during the international search (name of data base and, where practical, search terms used)

C. DOCUMENTS CONSIDERED TO BE RELEVANT

Category *	Citation of document, with indication, where appropriate, of the relevant passages	Relevant to claim No.
A	EP 0 476 439 A (BAYER AG) 25 March 1992 see claims 1-10 ---	1-12
A	EP 0 170 316 A (JANSSEN PHARMACEUTIA N.V.) 5 February 1986 cited in the application see claims 1-23 ---	1-12
A	EP 0 232 932 A (JANSSEN PHARMACEUTICA N.V.) 19 August 1987 cited in the application see claims 1-19 ---	1-12
A	EP 0 648 760 A (TAKEDA CHEMICAL INDUSTRIES) 19 April 1995 see claims 1-35 --- -/--	1-12

☒ Further documents are listed in the continuation of box C.

☒ Patent family members are listed in annex.

* Special categories of cited documents :

- "A" document defining the general state of the art which is not considered to be of particular relevance
- "E" earlier document but published on or after the international filing date
- "L" document which may throw doubts on priority claim(s) or which is cited to establish the publication date of another citation or other special reason (as specified)
- "O" document referring to an oral disclosure, use, exhibition or other means
- "P" document published prior to the international filing date but later than the priority date claimed

- "T" later document published after the international filing date or priority date and not in conflict with the application but cited to understand the principle or theory underlying the invention
- "X" document of particular relevance; the claimed invention cannot be considered novel or cannot be considered to involve an inventive step when the document is taken alone
- "Y" document of particular relevance; the claimed invention cannot be considered to involve an inventive step when the document is combined with one or more other such documents, such combination being obvious to a person skilled in the art.
- "&" document member of the same patent family

Date of the actual completion of the international search

3 November 1998

Date of mailing of the international search report

17/11/1998

Name and mailing address of the ISA

European Patent Office, P.B. 5818 Patentlaan 2
 NL - 2280 HV Rijswijk
 Tel. (+31-70) 340-2040, Tx. 31 651 epo nl,
 Fax: (+31-70) 340-3016

Authorized officer

Herz, C

INTERNATIONAL SEARCH REPORT

Intr. International Application No

PCT/EP 98/04191

C.(Continuation) DOCUMENTS CONSIDERED TO BE RELEVANT

Category	Citation of document, with indication, where appropriate, of the relevant passages	Relevant to claim No.
A	US 3 912 723 A (M. W. MILLER) 14 October 1975 see claims 1-21 -----	1-12
A	US 3 883 528 A (B. L. MYLARI) 13 May 1975 see claims 1-7 -----	1-12
A	DE 21 49 645 A (PFIZER INV.) 14 September 1972 see claims 1-3 -----	1-12

INTERNATIONAL SEARCH REPORT

Information on patent family members

International Application No

PCT/EP 98/04191

Patent document cited in search report		Publication date	Patent family member(s)	Publication date
EP 476439	A	25-03-1992	DE 4029534 A	19-03-1992
			DE 4120138 A	24-12-1992
			AU 639825 B	05-08-1993
			AU 8388691 A	26-03-1992
			CA 2051323 A	19-03-1992
			EG 19767 A	31-01-1996
			JP 5001047 A	08-01-1993
			US 5256631 A	26-10-1993
<hr/>				
EP 170316	A	05-02-1986	AU 579653 B	01-12-1988
			AU 4566485 A	06-02-1986
			BG 46599 A	15-01-1990
			CA 1244024 A	01-11-1988
			CA 1305479 A	21-07-1992
			CY 1656 A	14-05-1993
			DK 347585 A	02-02-1986
			EG 17589 A	30-12-1991
			FI 852958 A,B	02-02-1986
			GR 851837 A	02-12-1985
			HK 68092 A	18-09-1992
			IE 58076 B	30-06-1993
			JP 1732518 C	17-02-1993
			JP 4020435 B	02-04-1992
			JP 61043176 A	01-03-1986
			JP 1901539 C	27-01-1995
			JP 5017454 A	26-01-1993
			JP 6025177 B	06-04-1994
			LT 37093 R	15-06-1993
			LV 5027 A	10-06-1993
			MX 174156 B	25-04-1994
			PT 80903 B	11-11-1987
			SK 557685 A	04-09-1996
			SU 1443799 A	07-12-1988
			US 4631278 A	23-12-1986
			US 4778887 A	18-10-1988
<hr/>				
EP 232932	A	19-08-1987	AU 586731 B	20-07-1989
			AU 6813487 A	06-08-1987
			CA 1288772 A	10-09-1991
			CA 1302411 A	02-06-1992

INTERNATIONAL SEARCH REPORT

Information on patent family members

International Application No

PCT/EP 98/04191

Patent document cited in search report	Publication date	Patent family member(s)	Publication date
EP 232932 A		CN 1017049 B	17-06-1992
		DK 48187 A	31-07-1987
		DK 65193 A	04-06-1993
		FI 870421 A,B,	31-07-1987
		FI 923914 A	01-09-1992
		GR 3001392 T	11-09-1992
		IE 59510 B	09-03-1994
		JP 2098097 C	02-10-1996
		JP 8013806 B	14-02-1996
		JP 62190174 A	20-08-1987
		JP 2622510 B	18-06-1997
		JP 7285949 A	31-10-1995
		NO 175056 B	16-05-1994
		PT 84218 B	31-07-1989
		RU 2017736 C	15-08-1994
		SU 1595339 A	23-09-1990
		US 4767760 A	30-08-1988
		US 4912106 A	27-03-1990
		US 4952570 A	28-08-1990
EP 648760 A	19-04-1995	BR 9404113 A	27-06-1995
		CA 2118193 A	16-04-1995
		CN 1106394 A	09-08-1995
		JP 8143555 A	04-06-1996
US 3912723 A	14-10-1975	AR 195271 A	28-09-1973
		AR 196001 A	23-11-1973
		BE 773583 A	07-04-1972
		CA 992538 A	06-07-1976
		CH 566328 A	15-09-1975
		CH 563375 A	30-06-1975
		DE 2149645 A	14-09-1972
		FR 2110283 A	02-06-1972
		GB 1371907 A	30-10-1974
		NL 7113797 A	11-04-1972
		US 3905971 A	16-09-1975
US 3883528 A	13-05-1975	NONE	
DE 2149645 A	14-09-1972	AR 195271 A	28-09-1973

INTERNATIONAL SEARCH REPORT

Information on patent family members

International Application No

PCT/EP 98/04191

Patent document cited in search report	Publication date	Patent family member(s)	Publication date
DE 2149645 A		AR 196001 A	23-11-1973
		AT 325629 B	27-10-1975
		AT 866971 A	15-01-1975
		AU 3434771 A	12-04-1973
		BE 773583 A	07-04-1972
		CA 992538 A	06-07-1976
		CH 566328 A	15-09-1975
		CH 563375 A	30-06-1975
		FR 2110283 A	02-06-1972
		NL 7113797 A	11-04-1972
		ZA 7106613 A	28-06-1972
		GB 1371907 A	30-10-1974
		US 3905971 A	16-09-1975
		US 3912723 A	14-10-1975
